



Countdown to 2015

Maternal, Newborn & Child Survival

Accountability for Maternal, Newborn & Child Survival

An update on progress in priority countries

March 2012

Cover Photo: Mother with newborn baby, Ndosho maternity clinic, Goma, Democratic Republic of the Congo. WHO/Christopher Black.



THE LANCET



BILL & MELINDA GATES foundation



March 2012

Accountability for Maternal, Newborn & Child Survival

An update on progress in priority countries

www.countdown2015mnch.org



Countdown to 2015
Maternal, Newborn & Child Survival

© World Health Organization 2012

All rights reserved. Publications of the World Health Organization are available on the WHO web site (www.who.int).

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.

Contributors

This report has been developed by Countdown to 2015: Tracking Progress in Maternal, Newborn and Child Survival, in support of the Global Strategy for Women's and Children's Health and the follow-up to the report of the Commission on Information and Accountability for Women's and Children's Health, *Keeping Promises, Measuring Results*.

Countdown writing team: Jennifer Requejo (JHSPH/PMNCH), Jennifer Bryce (JHSPH), Adam Deixel (FCI), Cesar Victora (University of Pelotas)

Profile development team: Tessa Wardlaw (UNICEF), Archana Dwivedi (UNICEF), Holly Newby (UNICEF)

Production team: Virginia Taddoni (FCI), Adam Deixel (FCI), Lori McDougall (PMNCH), Dina El Husseiny (PMNCH), Jennifer Requejo (JHSPH/PMNCH)

Design: Virginia Taddoni and Sandy Zimmerman (FCI)

Printed in France by PAPRIKA

Acknowledgements

Countdown to 2015 would like to thank the following:

UNICEF/Statistics and Monitoring Section for use of global databases, preparation of country profiles and review of report text. Particular recognition goes to Priscilla Idele and Danielle Burke for their help with the HIV indicators.

DevInfo for its work on developing the profiles

The Countdown Scientific Review Group for its inputs in shaping the report: Cesar Victora (chair), Jennifer Bryce, Jennifer Requejo, Joy Lawn, Bernadette Daelmans, Holly Newby, Peter Berman, Andres de Francisco

The Countdown Leadership Sub-Group for its oversight of the report development process: Mickey Chopra (co-chair), Zulfiqar Bhutta (co-chair), Liz Mason, Carole Presern, Ann Starrs, Cesar Victora, Holly Newby, Jennifer Requejo



In this publication, Countdown to 2015

provides profiles for the countries where more than 95% of all maternal and child deaths occur. The profiles highlight how well each country is doing in increasing coverage of high-impact interventions that can save the lives of millions of women and children. The core indicators included in these profiles, selected in 2011 by the Commission on Information and Accountability for Women's and Children's Health, encompass key elements of the reproductive, maternal, newborn, and child health (RMNCH) continuum of care. A snapshot of progress on these core indicators, across the priority countries, reveals promising news as well as challenges that still remain to be addressed.



WHO/Pallava Bagla

COUNTDOWN TO 2015 IS A GLOBAL MOVEMENT OF ACADEMICS, GOVERNMENTS, INTERNATIONAL AGENCIES, HEALTH CARE PROFESSIONAL ORGANIZATIONS, DONORS, AND NON-GOVERNMENTAL ORGANIZATIONS, WITH THE LANCET AS A KEY PARTNER.

COUNTDOWN

- Uses country-specific data to track, stimulate, and support country progress towards achieving the health-related Millennium Development Goals (MDGs), particularly MDGs 4 (*reduce child mortality*) and 5 (*improve maternal health*)
- Focuses on coverage levels and trends for interventions proven to improve RMNCH, as well as critical determinants of coverage: health systems functionality, health policies, and financing
- Examines equity in coverage across different population groups within and across Countdown countries
- Uses these data to hold countries and their international partners accountable for progress in RMNCH
- Supports country-level Countdowns to promote evidence-based accountability

Since its first set of reports and events in 2005, Countdown has achieved global impact with its focus on accountability and use of available data to hold stakeholders to account for global and national action.

Countdown and the Accountability Agenda

In September 2010, at a UN General Assembly summit to assess progress on the MDGs, Secretary-General Ban Ki-moon launched the Global Strategy for Women's and Children's Health, an unprecedented plan to save the lives of 16 million women and children by 2015.¹ This was followed by the establishment of the Commission on Information and Accountability for Women's and Children's Health, charged with developing an accountability framework to monitor and track commitments made to the Global Strategy. In May 2011, the Commission released its report, *Keeping Promises, Measuring Results*. Drawing on advice from Countdown members and other technical experts, the report identified a set of core indicators which, taken together, enable stakeholders to track progress in improving coverage of interventions across the continuum of care. The report also urged that all coverage data be disaggregated by key equity considerations.² An independent Expert Review Group (iERG) was appointed by the Secretary-General in September 2011 to report annually on progress in implementing the Commission's recommendations regarding reporting, oversight, and accountability in 75 priority countries.³

Countdown is contributing significantly to this global accountability agenda, through the preparation of:

- Countdown profiles focused on the Commission indicators, updated every year with new data and results
- Special analyses to address accountability questions, and to inform the iERG
- Country-level Countdown processes that include national consultations, workshops, or publications, and utilize Countdown data and methodological approaches

The country profiles in this publication, customized to showcase the Commission indicators, are adapted from the full, two-page Countdown country profile, which Countdown produces on a roughly two-year cycle. Full country profiles will be included in Countdown's 2012 Report, to be published later in the year.

CORE INDICATORS IDENTIFIED BY THE COMMISSION ON INFORMATION AND ACCOUNTABILITY

- Maternal mortality ratio
- Under-5 child mortality, with the proportion of newborn deaths
- Children under five who are stunted
- Demand for family planning satisfied (met need for contraception)
- Antenatal care coverage (four or more visits)
- Antiretroviral prophylaxis among HIV-positive pregnant women to prevent mother-to-child transmission of HIV, and antiretroviral therapy for pregnant women who are treatment-eligible
- Skilled attendant at birth
- Postnatal care for mothers and babies within two days of birth
- Exclusive breastfeeding (< 6 months)
- Three doses of combined diphtheria-tetanus pertussis (DTP3) immunization coverage
- Antibiotic treatment for pneumonia

1. Up-to-date information on commitments to the Global Strategy is available at www.everywomaneverychild.org.

2. Further technical guidance on the Commission indicators is available from Countdown to 2015, Health Metrics Network, WHO. *Monitoring maternal, newborn and child health: understanding key progress indicators*. Geneva: WHO Document Production Services, 2011. ISBN 978 92 4 1502818 8.

3. Profiles are provided for 73 of the 75 high-burden countries. Profiles are not provided for South Sudan and post-cession Sudan because of lack of data availability at the time of publication.



WHO/Christopher Black

The Countdown country profile, highlighting the Commission indicators

The Countdown country profile presents in one place the best and latest evidence to enable an assessment of a country's progress in improving RMNCH. Most of the data for the Commission indicators come from household surveys: the two main surveys used to collect nationally-representative data for RMNCH are the USAID-supported Demographic and Health Surveys (DHS) and the UNICEF-supported Multiple Indicator Cluster Surveys (MICS).⁴

These country profiles reflect the data that are available for each country. Missing values for certain indicators, or estimates that are more than five years old, indicate an urgent need for concerted action to increase data collection efforts so that evidence is available for policy and program development.

Countdown addresses multiple Millennium Development Goals

- **MDG 4** to reduce child mortality
- **MDG 5** to improve maternal health
- **MDG 1** to eradicate extreme poverty and hunger, specifically by addressing nutrition with a focus on infant and young child feeding
- **MDG 6** to combat HIV/AIDS, malaria, and other diseases
- **MDG 7** to ensure environmental sustainability, through tracking improved access to safe water and improved sanitation

More information on the MDGs is available at:
www.un.org/millenniumgoals

4. More information about data sources is available at www.measuredhs.com and www.childinfo.org.

How to use the country profile

STEP 1

REVIEW THE DIFFERENT TYPES OF INFORMATION

The first step is to explore the range of data presented in the profile: demographic factors, mortality measures, coverage of evidence-based interventions, nutritional status measures, and measures of socioeconomic equity in coverage. Key questions to ask in reviewing the data include the following:

- Are trends in mortality and nutritional status moving in the right direction? Is the country on track to achieve the health MDGs?
- How high is coverage for the various interventions? Are trends moving in the right direction towards universal coverage? Are there gaps in coverage for specific interventions?
- How equitable is coverage? Are certain interventions particularly inaccessible for the poorest segment of the population?



Key population characteristics
These demographic indicators include proportion of newborn deaths among all under-5 mortality, a Commission indicator.

Continuum of care
Gaps in coverage along the continuum of care from pre-pregnancy and childbirth through childhood up to age 5 should serve as a call to action for a country to prioritize these interventions.

Equity in coverage
Socio-economic inequities in coverage highlight the need for concerted efforts to improve coverage among the poorest.

Impact: Under-5 child mortality rate and maternal mortality ratio
These charts display trends over time, reflecting progress towards reaching the MDG 4 and 5 targets.

Intervention coverage
These charts show most recent coverage levels and trends for selected RMNCH interventions.

Nutrition
Undernutrition contributes to at least 1/3 of all under-5 deaths, globally.

**STEP
2****IDENTIFY OPPORTUNITIES TO ACCELERATE PROGRESS**

The second step is to compare progress in different areas, focus on specific coverage gaps, and identify opportunities to address these gaps and accelerate progress in improving coverage and health outcomes. Questions to ask include the following:

- Are the coverage data consistent with the epidemiological situation? For example:
 - If stunting prevalence is high, are coverage levels low for the infant feeding indicators? Would a focus on early initiation of breastfeeding and exclusive breastfeeding help to drive progress on reducing stunting?
 - In priority countries for the elimination of mother-to-child transmission of HIV, are sufficient resources being targeted to preventing mother-to-child transmission?
 - If progress on reducing maternal mortality is lagging, and/or if newborn mortality is high, is this a reflection of low coverage of family planning, antenatal care (four or more visits), skilled attendance at birth, and postnatal care?
- Are there patterns in the coverage data that suggest clear action steps? For example, lower coverage for interventions involving treatment of an acute need (e.g., treatment of childhood diseases, childbirth services) than for interventions delivered routinely through outreach or scheduled in advance (e.g., vaccinations) suggests the need for measures to strengthen health systems, such as a greater policy focus on the training and equitable deployment of skilled health workers to increase access to care.
- Are gaps in coverage along the continuum of care suggestive of a call to action to prioritize specific interventions in future planning activities? For example, is access to labor, delivery, and immediate postnatal care being prioritized in countries with gaps in interventions delivered around the time of birth?

A snapshot of progress on the Commission indicators

This section summarizes the data presented in the country profiles included in this publication,⁵ providing an overall picture of progress and remaining challenges. Such summaries must be interpreted with care, because aggregate statistics can mask important information about progress or problems in individual countries.

IMPACT INDICATORS

These profiles show promising news on maternal mortality, though global progress is insufficient to achieve MDG 5. Thirty-five Countdown countries experienced a decline of 40% or more in maternal mortality between 1990 and 2008. Six countries are now on track to achieve MDG 5, and another 33 are making progress.⁶ However, progress is insufficient in 19 countries, and there has been no progress towards MDG 5 in another eight countries. Although maternal mortality has dropped in the majority of Countdown countries, 60% still have very high or high maternal mortality levels, indicating the need for continued focus on the provision of high-quality family planning, antenatal, delivery, and postnatal services.⁷

Under-5 mortality is declining! Among the Countdown countries, 24 are now on track to achieve MDG 4, up from 19

FIGURE 1
PROGRESS IN REDUCING MATERNAL MORTALITY (MDG 5)





FIGURE 2
PROGRESS IN REDUCING UNDER-5 MORTALITY (MDG 4)



5. Because of the secession in July 2011 of the Republic of South Sudan by the Republic of the Sudan, and its subsequent admission to the United Nations on 14 July 2011, disaggregated data for the Sudan and South Sudan as separate States are not yet available for most indicators. Aggregated data presented are for the Sudan pre-secession. Indicators for which Sudan (pre-secession) had available data during the time period 2006-2010 include: Demand for family planning satisfied, DTP3, Exclusive breastfeeding (<6 months), Skilled attendant at birth. This follows the convention used in UNICEF's State of the World's Children 2012 Report. The mortality and stunting data also include Sudan (pre-secession).

6. Key: Countries with MMR greater than or equal to 100 are categorized as: "on track" = 5.5% decline or more annually; "making progress" = declined between 2% and 5.5%; "insufficient progress" = less than 2% annually; "no progress" = an annual increase in MMR. Countries with MMR less than 100 in 1990 are not categorized. Source: WHO, UNICEF, UNFPA, World Bank, 2010. Trends in Maternal Mortality: 1990 to 2008. WHO.


7. WHO, UNICEF, UNFPA, World Bank, 2010. Trends in Maternal Mortality: 1990 to 2008. WHO.

at the time of the 2010 Countdown Report.⁸ The news is not all good, however, and about half of all under-5 deaths occur in only five Countdown countries: India, Nigeria, Democratic Republic of Congo, Pakistan, and China.

As under-5 mortality declines, the percentage of child deaths that are newborn deaths increases. The latest global estimates, as well as results from the 74 Countdown countries with data available, indicate that approximately 40% of under-5 deaths occur in the neonatal period.⁹ Countries must continue to prioritize child survival activities as well as concentrate on reducing newborn mortality.

Stunting is a critical indicator of progress in child survival, reflecting long-term exposure to poor health and nutrition, especially in the first two years of life. Children under the age of five around the world have the same growth potential, and prevalence of stunting above the 3% level expected in a well-nourished population indicates the need for remedial actions. All 61 Countdown countries with data available since 2006 have levels of stunting that require urgent attention. In the majority of these countries, more than one in three children is stunted. Addressing undernutrition through multi-sectoral programs must continue to be a major priority in these countries.

FIGURE 3
STUNTING PREVALENCE
 Percentage of children under five who are stunted, Countdown countries with data available, 2006-2010



8. Key: "On track" indicates that the under-5 mortality rate for 2010 is less than 40 per 1,000 or that it is 40 or more with an average annual rate of reduction of 4% or higher for 1990-2010; "insufficient progress" indicates that the under-5 mortality rate for 2010 is 40 or more with an average annual rate of reduction of 1% - 3.9% for 1990-2010; "no progress" indicates that the under-5 mortality rate for 2010 is 40 or more with an average annual rate of reduction of less than 1% for 1990 to 2010. Source: UNICEF, WHO, World Bank, United Nations Population Division/ DESA. 2011. *Levels & Trends in Child Mortality. Report 2011*. Estimates developed by the UN inter-agency group for child mortality estimation. UNICEF.

9. Source: UNICEF, WHO, World Bank, United Nations Population Division/DESA. 2011. *Levels & Trends in Child Mortality. Report 2011*. Estimates developed by the UN inter-agency group for child mortality estimation. UNICEF.

INDICATORS OF INTERVENTION COVERAGE

The Commission selected eight coverage indicators from among those already in use by Countdown and for tracking progress toward achieving the MDGs. Their choice was strategic, selecting a limited number of interventions along the continuum of care that are likely to be supported by the collection of consistent and timely data.

Figure 4 and Table 1 show the current coverage levels for the Commission indicators based on available data from the Countdown country profiles included in this publication. The HIV indicators selected by the Commission are presented separately below.

These coverage results demonstrate what is possible – for six of the interventions, at least one country has achieved a coverage level above 80%, and for the remaining two interventions, at least one country has achieved a coverage level of over 70%. The results also show, however, that substantial progress is still needed. The median coverage levels hover around 50%, with higher levels for DTP3 vaccine and lower levels for postnatal care for mothers, exclusive breastfeeding, and antibiotic treatment for pneumonia.

FIGURE 4
COVERAGE ACROSS THE CONTINUUM OF CARE
 Coverage levels for selected Commission indicators of intervention coverage, median and range for Countdown countries with data available, 2006-2010.




TABLE 1: A snapshot of coverage levels for select Commission indicators, Countdown countries with data available, 2006-2010.¹⁰

COMMISSION INDICATOR	NO. OF COUNTRIES WITH DATA	MEDIAN (%)	RANGE (%)
Demand for family planning satisfied (met need for contraception)	46	56	17 - 97
Antenatal care (four or more visits)	42	56	6 - 97
Skilled attendant at birth	65	57	18 - 100
Postnatal care (within two days)			
for mother	22	41	22 - 87
for baby	4	50	8 - 77
Exclusive breastfeeding (< 6 months)	57	37	1 - 74
DTP3 vaccine	74	85	33 - 99
Antibiotic treatment for pneumonia	41	38	3 - 88

Each country profile provides a summary graph showing whether coverage for a set of interventions is equitable across poorer and richer households. Analysis of the Countdown data shows that intervention coverage is substantially higher among women and children from richer households, but that inequalities in coverage vary by intervention. Interventions such as presence of a skilled attendant at birth are particularly inequitable, while interventions that do not require a functional health system, such as vaccines, are more equitable.¹¹

The Commission selected two HIV indicators, encouraging countries to increase provision of antiretroviral medicines to HIV-infected pregnant women in order to reduce the risk of transmission of HIV to their babies, and for their own health. These indicators are important measures of progress towards MDG 6. New reporting on coverage for the most effective antiretroviral drug regimens, as recommended by WHO, will now enable monitoring of country progress in scaling up these regimens. Table 2 shows coverage levels for the most effective regimens for the prevention of mother-to-child transmission (PMTCT) in the Countdown countries that are considered priority countries for the elimination of mother-to-child transmission. (Ethiopia, India, and Malawi are also priority countries, but they do not have disaggregated data for 2010, and are therefore not included.) The table shows a wide range in coverage levels, with three countries reporting coverage levels of 10% or less and six countries reaching over 75% of the eligible population in need.¹²

10. Source: UNICEF Global Databases, November 2011, based on Demographic and Health Surveys, Multiple Indicator Cluster Surveys and other national surveys.

11. Barros AJD, Ronemans C, Axelson H, Loaiza E, Bertali AD, Franca GV, Bryce J, Boerma T, Victora CG. Equity in maternal, newborn and child health interventions in Countdown to 2015: a retrospective review of survey data from 54 countries. *Lancet*, in press.

12. Source: WHO, UNAIDS, and UNICEF. 2011. *Global HIV/AIDS response. Epidemic update and health sector progress towards universal access*. Table 7.7, page. 156.

TABLE 2: Estimated antiretroviral coverage for the prevention of mother-to-child transmission using the most effective regimen, priority countries, 2010.

PRIORITY COUNTRY	PERCENT	UNCERTAINTY*
Congo, Democratic Republic of	1	(<1-1)
Chad	7	(5-9)
Nigeria	9	(7-10)
Angola	20	(15-28)
Burundi	36	(32-49)
Uganda	42	(36-51)
Kenya	43	(37-49)
Zimbabwe	46	(40-52)
Ghana	48	(40-57)
Mozambique	52	(44-62)
Cameroon	53	(43-65)
Tanzania, United Republic of	59	(52-68)
Côte d'Ivoire	66	(54-79)
Zambia	75	(67-85)
Lesotho	89	(77->95)
Botswana	>95	(>95->95)
South Africa	>95	(85->95)
Swaziland	>95	(88->95)

*The ranges around the levels of coverage are based on the uncertainty ranges around the estimates of need. Point estimates and ranges are given for countries with a generalized epidemic.

Coverage levels of antiretroviral therapy for HIV+ pregnant women who are treatment-eligible also vary substantially across the Countdown countries that are considered priorities for the elimination of mother-to-child transmission. Of the 17 priority countries with data for 2010, coverage values range from zero in Ghana to a high of 39% in Botswana and Chad.



WHO/Christopher Black

WHAT DO THE 2012 COUNTDOWN DATA TELL US?

- **There is important progress in reducing deaths among mothers and children, but efforts must be maintained and accelerated to achieve our goals in 2015 and beyond.**

Maternal and child survival gains must be protected and accelerated. Countries that are flagging must redouble their efforts; within countries—even those that are on track—population groups that are lagging behind must be identified and reached. Efforts targeted at reducing newborn deaths are urgent.

- **Undernutrition is a crisis that must be addressed now.**

Stunting prevalence is unacceptably high, even in countries that are making progress in reducing child mortality. Addressing child undernutrition should be a priority for all governments and their partners, especially among children under two years of age.

- **We must achieve universal coverage with proven RMNCH interventions.**

Only DTP3 coverage, with a median of 85% based on data from 74 countries, approaches an acceptable level of coverage. All other interventions are reaching fewer than 60% of women and children who need them. Inequities in intervention coverage must be corrected.

- **We can and must learn from country successes.**

The ranges in coverage are wide, indicating that for each intervention there are countries that are reaching much higher and much lower proportions of the population than the median. For example, although a median of 56% of women reported four or more antenatal care visits during their last pregnancy, this was true for only 6% of women in Somalia and for 97% of women in Swaziland. Countdown is analyzing the factors associated with differences in levels and trends of coverage in the priority countries as a basis for providing better guidance for governments and their partners.

Countdown country profiles: A basis for action

Information is a powerful force for change. Policies and programs must be developed based on evidence about what works and where improvements are needed. Governments, donors, and other stakeholders must be held accountable for fulfilling their commitments to improving RMNCH.

ACTIONS AT COUNTRY LEVEL

Countries can use the Countdown country profiles as a tool for action. Governments and their partners can conduct a “country Countdown” process, potentially involving conferences, reports, and the production of provincial or state-level Countdown profiles. These country Countdowns can:

- Stimulate policy and program improvements targeted at remedying coverage gaps and inequalities in coverage at national, provincial, district, and local levels
- Use evidence to advocate for more resources and greater political prioritization of the health of women and children

ACTIONS AT GLOBAL LEVEL

There is a role for each individual and every organization in increasing accountability for women’s and children’s health. The Countdown profiles offer a concise summary of the burden of maternal, newborn, and child mortality, and evidence on progress in efforts to scale-up proven interventions. Where the evidence shows that the level or focus of effort does not match the needs, decision makers and citizens have a responsibility to stand up and demand change.

Political will, combined with sound evidence, can change the world. Countdown stands ready to help.





Countdown to 2015

Maternal, Newborn & Child Survival

Country profiles

The information summarized in the profiles is intended to help policy makers and their partners assess progress, prioritize actions, and ensure accountability for national commitments to reduce maternal, newborn, and child mortality.

The following section contains profiles for 73 countries¹³ with high burdens of maternal and/or child mortality, including the 49 lowest-income countries covered by the Global Strategy for Women's and Children's Health. Profiles are included for the following countries:

Afghanistan	Gambia	Niger
Angola	Ghana	Nigeria
Azerbaijan	Guatemala	Pakistan
Bangladesh	Guinea	Papua New Guinea
Benin	Guinea-Bissau	Peru
Bolivia	Haiti	Philippines
Botswana	India	Rwanda
Brazil	Indonesia	Sao Tome and Principe
Burkina Faso	Iraq	Senegal
Burundi	Kenya	Sierra Leone
Cambodia	Korea, Democratic People's	Solomon Islands
Cameroon	Republic of	Somalia
Central African Republic	Kyrgyzstan	South Africa
Chad	Lao People's Democratic Republic	Swaziland
China	Lesotho	Tajikistan
Comoros	Liberia	Tanzania, United Republic of
Congo	Madagascar	Togo
Congo, Democratic Republic of the	Malawi	Turkmenistan
Côte d'Ivoire	Mali	Uganda
Djibouti	Mauritania	Uzbekistan
Egypt	Mexico	Vietnam
Equatorial Guinea	Morocco	Yemen
Eritrea	Mozambique	Zambia
Ethiopia	Myanmar	Zimbabwe
Gabon	Nepal	

13. There are 75 Countdown countries. Profiles for Sudan and South Sudan are not provided because of lack of available data at the time of publication.



DEMOGRAPHICS

Total population (000)	31,412	(2009)
Total under-five population (000)	5,546	(2009)
Births (000)	1,305	(2009)
Birth registration (%)	6	(2009)
Total under-five deaths (000)	191	(2009)
Neonatal deaths: % of all under-5 deaths	32	(2009)
Neonatal mortality rate (per 1000 live births)	45	(2009)
Infant mortality rate (per 1000 live births)	108	(2009)
Stillbirth rate (per 1000 total births)	29	(2009)
Total maternal deaths	18,000	(2009)
Lifetime risk of maternal death (1 in N)	11	(2009)
Total fertility rate (per woman)	6.3	(2009)
Adolescent birth rate (per 1000 women)	151	(2009)

Under-five mortality rate

Deaths per 1000 live births



Source: IHME 2010

Maternal mortality ratio

Deaths per 100,000 live births




Source: IMMR 2010

Note: MDG target established by Countdown in 2005

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See annexes for indicator details

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Skilled HIV-prepared women receiving ART for PMCT (Percent of total ART)

Percent HIV-prepared women receiving ART for PMCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 9 (2004)
Low birthweight incidence (moderate and severe, %) -

Early initiation of breastfeeding (within 1 hr of birth, %) -
Introduction of solid, semi-solid/soft foods (%) -
Vitamin A two dose coverage (%) 98 (2010)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2004 WHO reference population

Exclusive breastfeeding



Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	20,062	(2009)
Total under-five population (000)	8,376	(2009)
Births (000)	705	(2009)
Birth registration (%)	29	(2009)
Total under-five deaths (000)	121	(2009)
Neonatal deaths: % of all under-5 deaths	27	(2009)
Neonatal mortality rate (per 1000 live births)	41	(2009)
Infant mortality rate (per 1000 live births)	98	(2009)
Stillbirth rate (per 1000 total births)	25	(2009)
Total maternal deaths	4,830	(2009)
Lifetime risk of maternal death (1 in N)	29	(2009)
Total fertility rate (per woman)	5.4	(2009)
Adolescent birth rate (per 1000 women)	105	(2009)



MATERNAL AND NEWBORN HEALTH


Coverage along the continuum of care



Skilled attendant at delivery



Prevention of mother-to-child transmission of HIV




EQUITY

Socioeconomic inequities in coverage



CHILD HEALTH

Immunization




Pneumonia treatment



NUTRITION

Wasting prevalence (moderate and severe, %)	8	(2007)	Early initiation of breastfeeding (within 1 hr of birth, %)	-	(2007)
Low birthweight incidence (moderate and severe, %)	12	(2008)	Introduction of solid, semi-solid/soft foods (%)	-	(2007)
			Vitamin A two dose coverage (%)	28	(2008)

Underweight and stunting prevalence



Exclusive breastfeeding



Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

Note: Based on 2006 WHO reference population



DEMOGRAPHICS


Total population (000)	9,138	(2009)
Total under-five population (000)	795	(2009)
Births (000)	182	(2009)
Birth registration (%)	94	(2009)
Total under-five deaths (000)	9	(2009)
Neonatal deaths: % of all under-5 deaths	41	(2009)
Neonatal mortality rate (per 1000 live births)	39	(2009)
Infant mortality rate (per 1000 live births)	39	(2009)
Stillbirth rate (per 1000 total births)	12	(2009)
Total maternal deaths	63	(2009)
Lifetime risk of maternal death (1 in N)	1,200	(2009)
Total fertility rate (per woman)	2.1	(2010)
Adolescent birth rate (per 1000 women)	42	(2009)



Note: MDG target established by Countdown to 2015

MATERNAL AND NEWBORN HEALTH


Coverage along the continuum of care



* See annexes for indicator details


Skilled attendant at delivery

Percent live births attended by skilled health personnel




Prevention of mother-to-child transmission of HIV

Skilled HIV+ pregnant women receiving ART for their own health (% of total ARTs) 41 (2010)



EQUITY

Socioeconomic inequities in coverage




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	7	(2006)	Early initiation of breastfeeding (within 1 hr of birth, %)	32	(2006)
Low birthweight incidence (moderate and severe, %)	10	(2006)	Introduction of solid, semi-solid/soft foods (%)	83	(2006)
			Vitamin A two dose coverage (%)	89	(2010)


Underweight and stunting prevalence

Percent children <5 years who are underweight
Percent children <5 years who are stunted



Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	148,692	(1999)
Total under-five population (000)	24,707	(2008)
Births (000)	3,038	(2008)
Birth registration (%)	10	(2008)
Total under-five deaths (000)	140	(2008)
Neonatal deaths: % of all under-5 deaths	59	(2008)
Neonatal mortality rate (per 1000 live births)	27	(2008)
Infant mortality rate (per 1000 live births)	38	(2008)
Stillbirth rate (per 1000 total births)	36	(2008)
Total maternal deaths	12,000	(1990)
Lifetime risk of maternal death (1 in N)	101	(2008)
Total fertility rate (per woman)	2.2	(2008)
Adolescent birth rate (per 1000 women)	1.8	(2008)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ARTs for their own health (% of total HIV+)

Percent HIV+ pregnant women receiving ARTs for PMCT

Unertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2007

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

Percent of children immunized against measles

Percent of children immunized with 3 doses DTP

Percent of children immunized with 3 doses Hib




Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider

Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)

Low birthweight incidence (moderate and severe, %)

Early initiation of breastfeeding (within 1 hr of birth, %)


Introduction of solid, semi-solid/soft foods (%)

Vitamin A two dose coverage (%)

Underweight and stunting prevalence

Percent children <5 years who are underweight


Percent children <5 years who are stunted



Note: Based on 2000 WHO reference population

Exclusive breastfeeding



Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	8,850	(2009)
Total under-five population (000)	1,506	(2009)
Births (000)	350	(2009)
Birth registration (%)	60	(2009)
Total under-five deaths (000)	39	(2009)
Neonatal deaths: % of all under-5 deaths	29	(2009)
Neonatal mortality rate (per 1000 live births)	32	(2009)
Infant mortality rate (per 1000 live births)	73	(2009)
Skilled birth rate (per 1000 total births)	24	(2009)
Total maternal deaths	1,430	(2009)
Lifetime risk of maternal death (1 in N)	48	(2009)
Total fertility rate (per woman)	5.3	(2010)
Adolescent birth rate (per 1000 women)	114	(2009)




MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Skilled attendant at delivery




Prevention of mother-to-child transmission of HIV




EQUITY

Socioeconomic inequities in coverage




CHILD HEALTH

Immunization




Pneumonia treatment



NUTRITION

Wasting prevalence (moderate and severe, %)	8	(2008)	Early initiation of breastfeeding (within 1 hr of birth, %)	33	(2008)
Low birthweight incidence (moderate and severe, %)	15	(2008)	Introduction of solid, semi-solid/soft foods (%)	-	-
			Vitamin A two dose coverage (%)	100	(2008)

Underweight and stunting prevalence



Exclusive breastfeeding






DEMOGRAPHICS

Total population (000)	9,930	(2008)
Total under-five population (000)	1,235	(2008)
Births (000)	263	(2008)
Birth registration (%)	34	(2008)
Total under-five deaths (000)	14	(2008)
Neonatal deaths: % of all under-5 deaths	44	(2008)
Neonatal mortality rate (per 1000 live births)	23	(2008)
Infant mortality rate (per 1000 live births)	42	(2008)
Stillbirth rate (per 1000 total births)	17	(2008)
Total maternal deaths	470	(2008)
Lifetime risk of maternal death (1 in N)	150	(2008)
Total fertility rate (per woman)	3.3	(2008)
Adolescent birth rate (per 1000 women)	89	(2008)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV


Eligible HIV+ pregnant women receiving ART for PMCT (% of total HIV+)



Source: UNICEF/UNAIDS/WHO

EQUITY


Socioeconomic inequities in coverage



DHS 2008
Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.


CHILD HEALTH

Immunization



Source: WHO/UNICEF


Pneumonia treatment



NUTRITION

Wasting prevalence (moderate and severe, %)	1	(2008)
Low birthweight incidence (moderate and severe, %)	6	(2008)
Early initiation of breastfeeding (within 1 hr of birth, %)	64	(2008)
Introduction of solid, semi-solid/soft foods (%)	83	(2008)
Vitamin A two dose coverage (%)	24	(2008)

Underweight and stunting prevalence



Note: Based on 2008 WHO reference population



Exclusive breastfeeding






DEMOGRAPHICS

Total population (000)	2,037	(2009)
Total under-five population (000)	225	(2009)
Births (000)	47	(2009)
Birth registration (%)	72	(2009)
Total under-five deaths (000)	2	(2009)
Neonatal deaths: % of all under-5 deaths	40	(2009)
Neonatal mortality rate (per 1000 live births)	19	(2009)
Infant mortality rate (per 1000 live births)	36	(2009)
Stillbirth rate (per 1000 total births)	36	(2009)
Total maternal deaths	91	(2009)
Lifetime risk of maternal death (1 in N)	180	(2009)
Total fertility rate (per woman)	2.8	(2009)
Adolescent birth rate (per 1000 women)	51	(2009)




MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Skilled attendant at delivery



Prevention of mother-to-child transmission of HIV



EQUITY

Socioeconomic inequities in coverage




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on care indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization




Pneumonia treatment



NUTRITION

Wasting prevalence (moderate and severe, %)	7	(2006)	Early initiation of breastfeeding (within 1 hr of birth, %)	40	(2008)
Low birthweight incidence (moderate and severe, %)	13	(2007)	Introduction of solid, semi-solid/soft foods (%)	-	-
			Vitamin A two dose coverage (%)	81	(2008)

Underweight and stunting prevalence



Exclusive breastfeeding






DEMOGRAPHICS

Total population (000)	254,946	(2009)
Total under-five population (000)	15,136	(2009)
Births (000)	3,023	(2009)
Birth registration (%)	91	(2009)
Total under-five deaths (000)	55	(2009)
Neonatal deaths: % of all under-5 deaths	65	(2009)
Neonatal mortality rate (per 1000 live births)	12	(2009)
Infant mortality rate (per 1000 live births)	17	(2009)
SBIR* birth rate (per 1000 total births)	36	(2009)
Total maternal deaths	1,800	(2009)
Lifetime risk of maternal death (1 in N)	960	(2009)
Total fertility rate (per woman)	1.8	(2009)
Adolescent birth rate (per 1000 women)	77	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See annex table for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Source: UNICEF/UNAIDS/WHO

Prevention of mother-to-child transmission of HIV

Highly HIV+ pregnant women receiving ART for their own health (% of total HIV+)

Percent HIV+ pregnant women receiving ART for PMCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)	2	(2006)	Early initiation of breastfeeding (within 1 hr of birth, %)	33	(2006)
Low birthweight incidence (moderate and severe, %)	8	(2006)	Introduction of solid, semi-solid/soft foods (%)	-	-
			Vitamin A two dose coverage (%)	-	-

Underweight and stunting prevalence

Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding



Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	16,059	(2009)
Total under-five population (000)	2,950	(2009)
Births (000)	713	(2009)
Birth registration (%)	64	(2009)
Total under-five deaths (000)	120	(2009)
Neonatal deaths (% of all under-5 deaths)	22	(2009)
Neonatal mortality rate (per 1000 live births)	30	(2009)
Infant mortality rate (per 1000 live births)	98	(2009)
Stillbirth rate (per 1000 total births)	26	(2009)
Total maternal deaths	4,000	(2009)
Lifetime risk of maternal death (1 in N)	28	(2009)
Total fertility rate (per woman)	5.9	(2009)
Adolescent birth rate (per 1000 women)	128	(2009)



MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Skilled attendant at delivery




Prevention of mother-to-child transmission of HIV




EQUITY

Socioeconomic inequities in coverage




CHILD HEALTH

Immunization




Pneumonia treatment



NUTRITION

Wasting prevalence (moderate and severe, %)	11	(2008)
Low birthweight incidence (moderate and severe, %)	18	(2008)
Early initiation of breastfeeding (within 1 hr of birth, %)	20	(2008)
Introduction of solid, semi-solid/soft foods (%)	-	-
Vitamin A two dose coverage (%)	100	(2008)

Underweight and stunting prevalence





Exclusive breastfeeding






DEMOGRAPHICS

Total population (000)	8,363	(2009)
Total under-five population (000)	1,185	(2009)
Births (000)	283	(2009)
Birth registration (%)	60	(2008)
Total under-five deaths (000)	39	(2008)
Neonatal deaths (% of all under-5 deaths)	31	(2008)
Neonatal mortality rate (per 1000 live births)	42	(2008)
Infant mortality rate (per 1000 live births)	88	(2008)
Skilled birth rate (per 1000 total births)	28	(2008)
Total maternal deaths	2,700	(2008)
Lifetime risk of maternal death (1 in N)	29	(2008)
Total fertility rate (per woman)	4.3	(2008)
Adolescent birth rate (per 1000 women)	30	(2003)



MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS
* See annexes for indicator details


Skilled attendant at delivery

Percent live births attended by skilled health personnel




Prevention of mother-to-child transmission of HIV

Skilled HIV+ pregnant women receiving ART for their own health (% of total ART)



EQUITY


Socioeconomic inequities in coverage



MICS 2005
Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH


Immunization



Source: WHO/UNICEF

Pneumonia treatment


Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)	6	(2008)
Low birthweight incidence (moderate and severe, %)	11	(2008)
Early initiation of breastfeeding (within 1 hr of birth, %)	-	-
Introduction of solid, semi-solid/soft foods (%)	-	-
Vitamin A two dose coverage (%)	71	(2008)

Underweight and stunting prevalence



Note: Based on 2006 WHO reference population

Exclusive breastfeeding






DEMOGRAPHICS

Total population (000)	14,186	(2009)
Total under-five population (000)	1,492	(2009)
Births (000)	318	(2009)
Birth registration (%)	62	(2009)
Total under-five deaths (000)	16	(2009)
Neonatal deaths: % of all under-5 deaths	42	(2009)
Neonatal mortality rate (per 1000 live births)	22	(2009)
Infant mortality rate (per 1000 live births)	43	(2009)
Stillbirth rate (per 1000 total births)	16	(2009)
Total maternal deaths	1,100	(2009)
Lifetime risk of maternal death (1 in N)	101	(2009)
Total fertility rate (per woman)	2.6	(2009)
Adolescent birth rate (per 1000 women)	52	(2009)

Under-five mortality rate

Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV


Eligible HIV+ pregnant women receiving ARTs for PMCT (% of total HIV+)



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage




DHS 2010

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider




Source: DHS, MICS, Other NS

NUTRITION

Wasting prevalence (moderate and severe, %)	11	(2008)
Low birthweight incidence (moderate and severe, %)	9	(2008)
Early initiation of breastfeeding (within 1 hr of birth, %)	65	(2008)
Introduction of solid, semi-solid/soft foods (%)	-	(2008)
Vitamin A two dose coverage (%)	88	(2008)

Underweight and stunting prevalence



Note: Based on 2000 WHO reference population



Exclusive breastfeeding






DEMOGRAPHICS

Total population (000)	19,599	(2009)
Total under-five population (000)	3,050	(2009)
Births (000)	730	(2009)
Birth registration (%)	70	(2009)
Total under-five deaths (000)	93	(2009)
Neonatal deaths: % of all under-5 deaths	26	(2009)
Neonatal mortality rate (per 1000 live births)	34	(2009)
Infant mortality rate (per 1000 live births)	94	(2009)
Fertility rate (per 1000 total births)	26	(2009)
Total maternal deaths	4,200	(2009)
Lifetime risk of maternal death (1 in N)	85	(2009)
Total fertility rate (per woman)	4.5	(2009)
Adolescent birth rate (per 1000 women)	141	(2009)



MATERNAL AND NEWBORN HEALTH


Coverage along the continuum of care



Skilled attendant at delivery




Prevention of mother-to-child transmission of HIV




EQUITY

Socioeconomic inequities in coverage




CHILD HEALTH

Immunization



Pneumonia treatment



NUTRITION

Wasting prevalence (moderate and severe, %)	7	(2006)	Early initiation of breastfeeding (within 1 hr of birth, %)	20	(2006)
Low birthweight incidence (moderate and severe, %)	11	(2008)	Introduction of solid, semi-solid/soft foods (%)	63	(2006)
			Vitamin A two dose coverage (%)	89	(2010)

Underweight and stunting prevalence



Exclusive breastfeeding






DEMOGRAPHICS

Total population (000)	4,401	(2008)
Total under-five population (000)	951	(2008)
Births (000)	154	(2008)
Birth registration (%)	40	(2008)
Total under-five deaths (000)	23	(2008)
Neonatal deaths: % of all under-5 deaths	30	(2008)
Neonatal mortality rate (per 1000 live births)	42	(2008)
Infant mortality rate (per 1000 live births)	106	(2008)
Stillbirth rate (per 1000 total births)	24	(2008)
Total maternal deaths	1,300	(2008)
Lifetime risk of maternal death (1 in N)	37	(2008)
Total fertility rate (per woman)	4.6	(2008)
Adolescent birth rate (per 1000 women)	133	(2008)

Under-five mortality rate

Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total HIV+)

Percent HIV+ pregnant women receiving ART for PMCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




MICS 2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	12	(2006)
Low birthweight incidence (moderate and severe, %)	13	(2006)
Early initiation of breastfeeding (within 1 hr of birth, %)	39	(2006)
Introduction of solid, semi-solid/soft foods (%)	56	(2006)
Vitamin A two dose coverage (%)	0	(2006)

Underweight and stunting prevalence

Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding



Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	11,227	(2008)
Total under-five population (000)	2,036	(2008)
Births (000)	508	(2008)
Birth registration (%)	9	(2008)
Total under-five deaths (000)	80	(2008)
Neonatal deaths: % of all under-5 deaths	26	(2008)
Neonatal mortality rate (per 1000 live births)	41	(2008)
Infant mortality rate (per 1000 live births)	99	(2008)
Stillbirth rate (per 1000 total births)	29	(2008)
Total maternal deaths	5,000	(2008)
Lifetime risk of maternal death (1 in N)	50	(2008)
Total fertility rate (per woman)	6.0	(2008)
Adolescent birth rate (per 1000 women)	198	(2008)



MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Skilled attendant at delivery




Prevention of mother-to-child transmission of HIV




EQUITY

Socioeconomic inequities in coverage




CHILD HEALTH

Immunization




Pneumonia treatment



NUTRITION

Wasting prevalence (moderate and severe, %)	36	(2008)
Low birthweight incidence (moderate and severe, %)	22	(2008)
Early initiation of breastfeeding (within 1 hr of birth, %)	34	(2008)
Introduction of solid, semi-solid/soft foods (%)	36	(2010)
Vitamin A two dose coverage (%)	68	(2010)

Underweight and stunting prevalence




Exclusive breastfeeding






DEMOGRAPHICS

Total population (000)	1,341,805	(2009)
Total under-five population (000)	81,596	(2009)
Births (000)	16,489	(2009)
Birth registration (%)	-	-
Total under-five deaths (000)	123	(2009)
Neonatal deaths: % of all under-five deaths	58	(2009)
Neonatal mortality rate (per 1000 live births)	11	(2009)
Infant mortality rate (per 1000 live births)	26	(2009)
SBIR rate (per 1000 total births)	30	(2009)
Total maternal deaths	6,900	(2009)
Lifetime risk of maternal death (1 in N)	1,500	(2009)
Total fertility rate (per woman)	1.6	(2010)
Adolescent birth rate (per 1000 women)	6	(2009)




MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Skilled attendant at delivery




Prevention of mother-to-child transmission of HIV



EQUITY

Socioeconomic inequities in coverage



CHILD HEALTH

Immunization




Pneumonia treatment



NUTRITION

Wasting prevalence (moderate and severe, %)	3	(2008)	Early initiation of breastfeeding (within 1 hr of birth, %)	41	(2008)
Low birthweight incidence (moderate and severe, %)	3	(2008)	Introduction of solid, semi-solid/soft foods (%)	-	-
			Vitamin A two dose coverage (%)	-	-

Underweight and stunting prevalence



Exclusive breastfeeding





Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.




DEMOGRAPHICS

Total population (000)	785	(2009)
Total under-five population (000)	222	(2009)
Births (000)	28	(2009)
Birth registration (%)	63	(2009)
Total under-five deaths (000)	2	(2009)
Neonatal deaths (% of all under-5 deaths)	39	(2009)
Neonatal mortality rate (per 1000 live births)	32	(2009)
Infant mortality rate (per 1000 live births)	58	(2009)
Skilled birth rate (per 1000 total births)	27	(2009)
Total maternal deaths	72	(2009)
Lifetime risk of maternal death (1 in N)	71	(2009)
Total fertility rate (per woman)	4.9	(2009)
Adolescent birth rate (per 1000 women)	95	(2009)




MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Skilled attendant at delivery

Percent live births attended by skilled health personnel




Prevention of mother-to-child transmission of HIV

Skilled HIV-prepared women receiving ART for their own health (% of total ART)




EQUITY

Socioeconomic inequities in coverage




CHILD HEALTH

Immunization



Pneumonia treatment



NUTRITION

Wasting prevalence (moderate and severe, %)	-	Early initiation of breastfeeding (within 1 hr of birth, %)	25	(2006)	
Low birthweight incidence (moderate and severe, %)	25	(2008)	Introduction of solid, semi-solid/soft foods (%)	-	
			Vitamin A two dose coverage (%)	18	(2008)

Underweight and stunting prevalence



Exclusive breastfeeding






DEMOGRAPHICS

Total population (000)	4,048	(2009)
Total under-five population (000)	923	(2009)
Births (000)	142	(2009)
Birth registration (%)	81	(2009)
Total under-five deaths (000)	13	(2009)
Neonatal deaths: % of all under-5 deaths	32	(2009)
Neonatal mortality rate (per 1000 live births)	29	(2009)
Infant mortality rate (per 1000 live births)	64	(2009)
Stillbirth rate (per 1000 total births)	26	(2009)
Total maternal deaths	720	(2009)
Lifetime risk of maternal death (1 in N)	89	(2009)
Total fertility rate (per woman)	4.5	(2009)
Adolescent birth rate (per 1000 women)	112	(2009)

Under-five mortality rate

Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: WHO 2009

Note: MDG target established by Countdown to 2015

MATERNAL AND NEWBORN HEALTH


Coverage along the continuum of care



Source: DHS, MICS, Other NS
* See annex tables for indicator details

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Skilled HIV-prepared women receiving ART for PMCT
Share of total ART (%)


Percent HIV-prepared women receiving ART for PMCT
Uncertainty range around the estimate



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage



Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)	8	(2008)
Low birthweight incidence (moderate and severe, %)	13	(2008)
Early initiation of breastfeeding (within 1 hr of birth, %)	39	(2008)
Introduction of solid, semi-solid/soft foods (%)	-	-
Vitamin A two dose coverage (%)	84	(2008)

Underweight and stunting prevalence

Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding



Percent infants <6 months exclusively breastfed





DEMOGRAPHICS


Total population (000)	65,956	(2010)
Total under-five population (000)	11,648	(2010)
Births (000)	2,879	(2010)
Birth registration (%)	28	(2010)
Total under-five deaths (000)	465	(2010)
Neonatal deaths, % of all under-five deaths	29	(2010)
Neonatal mortality rate (per 1000 live births)	40	(2010)
Infant mortality rate (per 1000 live births)	112	(2010)
Stillbirth rate (per 1000 total births)	29	(2010)
Total maternal deaths	10,031	(2010)
Lifetime risk of maternal death (1 in N)	34	(2010)
Total fertility rate (per woman)	5.8	(2010)
Adolescent birth rate (per 1000 women)	15	(2010)



Note: MDG target established by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



* See annexure for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Skilled HIV+ pregnant women receiving at least one health (% of total HIV+)


■ Percent HIV+ pregnant women receiving ART for PMCT
■ Uncertainty range around the estimate



EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH


Immunization

■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	9	(2010)
Low birthweight incidence (moderate and severe, %)	10	(2010)
Early initiation of breastfeeding (within 1 hr of birth, %)	43	(2010)
Introduction of solid, semi-solid/soft foods (%)	57	(2010)
Vitamin A two dose coverage (%)	83	(2010)

Underweight and stunting prevalence

■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	18,736	(2008)
Total under-five population (000)	2,940	(2008)
Births (000)	673	(2008)
Birth registration (%)	55	(2008)
Total under-five deaths (000)	80	(2008)
Neonatal deaths: % of all under-5 deaths	34	(2008)
Neonatal mortality rate (per 1000 live births)	41	(2008)
Infant mortality rate (per 1000 live births)	96	(2008)
Stillbirth rate (per 1000 total births)	27	(2008)
Total maternal deaths	3,400	(2008)
Lifetime risk of maternal death (1 in N)	44	(2008)
Total fertility rate (per woman)	4.4	(2008)
Adolescent birth rate (per 1000 women)	111	(2008)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ARTs for PMCT (% of total HIV+ live births) 17 (2010)


Percent HIV+ pregnant women receiving ARTs for PMCT



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage



MICS 2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics



Source: DHS, MICS, Other NS

NUTRITION

Wasting prevalence (moderate and severe, %)	9	(2008)
Low birthweight incidence (moderate and severe, %)	17	(2008)
Early initiation of breastfeeding (within 1 hr of birth, %)	25	(2008)
Introduction of solid, semi-solid/soft foods (%)	81	(2008)
Vitamin A two dose coverage (%)	100	(2008)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2008 WHO reference population

Exclusive breastfeeding



Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	688	(2008)
Total under-five population (000)	113	(2008)
Births (000)	26	(2008)
Birth registration (%)	89	(2008)
Total under-five deaths (000)	2	(2008)
Neonatal deaths: % of all under-5 deaths	38	(2008)
Neonatal mortality rate (per 1000 live births)	34	(2008)
Infant mortality rate (per 1000 live births)	73	(2008)
Skilled birth rate (per 1000 total births)	94	(2008)
Total maternal deaths	73	(2008)
Lifetime risk of maternal death (1 in N)	68	(2008)
Total fertility rate (per woman)	4.8	(2008)
Adolescent birth rate (per 1000 women)	27	(2008)




MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Skilled attendant at delivery



Prevention of mother-to-child transmission of HIV




EQUITY

Socioeconomic inequities in coverage




CHILD HEALTH

Immunization




Pneumonia treatment



NUTRITION

Wasting prevalence (moderate and severe, %)	30	(2008)
Low birthweight incidence (moderate and severe, %)	30	(2008)
Early initiation of breastfeeding (within 1 hr of birth, %)	67	(2008)
Introduction of solid, semi-solid/soft foods (%)	35	(2008)
Vitamin A two dose coverage (%)	95	(2008)

Underweight and stunting prevalence



Exclusive breastfeeding



Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

Note: Based on 2008 WHO reference population




DEMOGRAPHICS

Total population (000)	81,321	(2008)
Total under-five population (000)	9,008	(2008)
Births (000)	1,881	(2008)
Birth registration (%)	99	(2008)
Total under-five deaths (000)	41	(2008)
Neonatal deaths: % of all under-5 deaths	44	(2008)
Neonatal mortality rate (per 1000 live births)	9	(2008)
Infant mortality rate (per 1000 live births)	19	(2008)
Stillbirth rate (per 1000 total births)	11	(2008)
Total maternal deaths	1,600	(2008)
Lifetime risk of maternal death (1 in N)	380	(2008)
Total fertility rate (per woman)	2.7	(2008)
Adolescent birth rate (per 1000 women)	30	(2008)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Source: UNICEF/UNAIDS/WHO

Prevention of mother-to-child transmission of HIV

Highly HIV+ pregnant women receiving ART for their own health (% of total HIV+)

Percent HIV+ pregnant women receiving ART for PMCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2008

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)	8	(2008)	Early initiation of breastfeeding (within 1 hr of birth, %)	43	(2008)
Low birthweight incidence (moderate and severe, %)	13	(2008)	Introduction of solid, semi-solid/soft foods (%)	70	(2008)
			Vitamin A two dose coverage (%)	68	(2008)

Underweight and stunting prevalence

Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2008 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	700	(2008)
Total under-five population (000)	307	(2008)
Births (000)	26	(2008)
Birth registration (%)	32	(2008)
Total under-five deaths (000)	3	(2008)
Neonatal deaths: % of all under-5 deaths	31	(2008)
Neonatal mortality rate (per 1000 live births)	35	(2008)
Infant mortality rate (per 1000 live births)	81	(2008)
Stillbirth rate (per 1000 total births)	17	(2008)
Total maternal deaths	20	(2008)
Lifetime risk of maternal death (1 in N)	73	(2008)
Total fertility rate (per woman)	5.2	(2008)
Adolescent birth rate (per 1000 women)	128	(2008)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annex table for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for PMCT as a % of total HIV+ (2010)



Source: UNICEF/UNAIDS/WHO

EQUITY


Socioeconomic inequities in coverage



Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH


Immunization



Source: WHO/UNICEF

Pneumonia treatment


Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)	9	(2008)	Early initiation of breastfeeding (within 1 hr of birth, %)	-	(2008)
Low birthweight incidence (moderate and severe, %)	13	(2008)	Introduction of solid, semi-solid/soft foods (%)	-	(2008)
			Vitamin A two dose coverage (%)	0	(2008)

Underweight and stunting prevalence



Note: Based on 2000 WHO reference population



Exclusive breastfeeding





DEMOGRAPHICS


Total population (000)	5,254	(2009)
Total under-five population (000)	861	(2009)
Births (000)	178	(2009)
Birth registration (%)	-	(2009)
Total under-five deaths (000)	11	(2009)
Neonatal deaths: % of all under-5 deaths	31	(2009)
Neonatal mortality rate (per 1000 live births)	38	(2009)
Infant mortality rate (per 1000 live births)	42	(2009)
Skilled birth rate (per 1000 total births)	21	(2009)
Total maternal deaths	520	(2009)
Lifetime risk of maternal death (1 in N)	72	(2009)
Total fertility rate (per woman)	4.5	(2009)
Adolescent birth rate (per 1000 women)	85	(2009)



Note: MDG target established by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Skilled HIV-prepared women receiving ART for their own health (% of total ART)


Percent HIV-prepared women receiving ART for PMCT
Uncertainty range around the estimate



EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	25	(2008)	Early initiation of breastfeeding (within 1 hr of birth, %)	78	(2008)
Low birthweight incidence (moderate and severe, %)	14	(2002)	Introduction of solid, semi-solid/soft foods (%)	-	-
			Vitamin A two dose coverage (%)	44	(2008)


Underweight and stunting prevalence

Percent children <5 years who are underweight
Percent children <5 years who are stunted



Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	82,950	(2009)
Total under-five population (000)	11,942	(2009)
Births (000)	2,653	(2009)
Birth registration (%)	7	(2009)
Total under-five deaths (000)	271	(2009)
Neonatal deaths: % of all under-5 deaths	34	(2009)
Neonatal mortality rate (per 1000 live births)	35	(2009)
Infant mortality rate (per 1000 live births)	68	(2009)
SBIR rate (per 1000 total births)	26	(2009)
Total maternal deaths	14,000	(2009)
Lifetime risk of maternal death (1 in N)	40	(2009)
Total fertility rate (per woman)	4.2	(2009)
Adolescent birth rate (per 1000 women)	109	(2009)

Under-five mortality rate

Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Highly HIV+ pregnant women receiving ARTs for PMCT (% of total HIV+)

14 (2010)




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2005

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	12	(2008)	Early initiation of breastfeeding (within 1 hr of birth, %)	69	(2008)
Low birthweight incidence (moderate and severe, %)	20	(2008)	Introduction of solid, semi-solid/soft foods (%)	-	-
			Vitamin A two dose coverage (%)	84	(2008)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2000 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	1,505	(2009)
Total under-five population (000)	185	(2009)
Births (000)	41	(2009)
Birth registration (%)	89	(2009)
Total under-five deaths (000)	3	(2009)
Neonatal deaths: % of all under-5 deaths	37	(2009)
Neonatal mortality rate (per 1000 live births)	26	(2009)
Infant mortality rate (per 1000 live births)	54	(2009)
SBIR* birth rate (per 1000 total births)	17	(2009)
Total maternal deaths	100	(2009)
Lifetime risk of maternal death (1 in N)	110	(2009)
Total fertility rate (per woman)	3.3	(2009)
Adolescent birth rate (per 1000 women)	144	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Source: DHS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for PMCT (% of total HIV+)

Percent HIV+ pregnant women receiving ART for PMCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2000

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics




Source: DHS

NUTRITION

Wasting prevalence (moderate and severe, %)	4	(2008)	Early initiation of breastfeeding (within 1 hr of birth, %)	71	(2008)
Low birthweight incidence (moderate and severe, %)	14	(2008)	Introduction of solid, semi-solid/soft foods (%)	-	-
			Vitamin A two dose coverage (%)	0	(2008)

Underweight and stunting prevalence

Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2000 WHO reference population

Exclusive breastfeeding



Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	1,726	(2009)
Total under-five population (000)	267	(2009)
Births (000)	56	(2009)
Birth registration (%)	50	(2008-2009)
Total under-five deaths (000)	6	(2009)
Neonatal deaths (% of all under-5 deaths)	34	(2009)
Neonatal mortality rate (per 1000 live births)	31	(2009)
Infant mortality rate (per 1000 live births)	57	(2009)
Stillbirth rate (per 1000 total births)	26	(2009)
Total maternal deaths	250	(2009)
Lifetime risk of maternal death (1 in N)	69	(2009)
Total fertility rate (per woman)	4.9	(2009)
Adolescent birth rate (per 1000 women)	104	(2009)




MATERNAL AND NEWBORN HEALTH


Coverage along the continuum of care



Skilled attendant at delivery




Prevention of mother-to-child transmission of HIV




EQUITY

Socioeconomic inequities in coverage




CHILD HEALTH

Immunization




Pneumonia treatment



NUTRITION

Wasting prevalence (moderate and severe, %)	7 (2005-2006)	Early initiation of breastfeeding (within 1 hr of birth, %)	48 (2009-2006)
Low birthweight incidence (moderate and severe, %)	20 (2005-2006)	Introduction of solid, semi-solid/soft foods (%)	-
		Vitamin A two dose coverage (%)	100 (2006)

Underweight and stunting prevalence




Exclusive breastfeeding





DEMOGRAPHICS


Total population (000)	24,982	(2009)
Total under-five population (000)	3,533	(2009)
Births (000)	770	(2009)
Birth registration (%)	71	(2009)
Total under-five deaths (000)	57	(2009)
Neonatal deaths (% of all under-5 deaths)	38	(2009)
Neonatal mortality rate (per 1000 live births)	26	(2009)
Infant mortality rate (per 1000 live births)	50	(2009)
Skilled birth rate (per 1000 total births)	22	(2009)
Total maternal deaths	2,600	(2009)
Lifetime risk of maternal death (1 in N)	66	(2009)
Total fertility rate (per woman)	4.2	(2009)
Adolescent birth rate (per 1000 women)	70	(2009)



Note: MDG target established by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



* See annexes for indicator details


Skilled attendant at delivery

Percent live births attended by skilled health personnel




Prevention of mother-to-child transmission of HIV

Skilled HIV+ pregnant women receiving ART to reduce health (% of total ART)



EQUITY

Socioeconomic inequities in coverage




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on care indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization



Pneumonia treatment


Percent children <5 years with suspected pneumonia taken to appropriate health provider



NUTRITION

Wasting prevalence (moderate and severe, %)	9	(2008)
Low birthweight incidence (moderate and severe, %)	13	(2008)
Early initiation of breastfeeding (within 1 hr of birth, %)	53	(2008)
Introduction of solid, semi-solid/soft foods (%)	76	(2008)
Vitamin A two dose coverage (%)	83	(2008)

Underweight and stunting prevalence



Exclusive breastfeeding






DEMOGRAPHICS

Total population (000)	14,389	(1999)
Total under-five population (000)	2,167	(1999)
Births (000)	427	(1999)
Birth registration (%)	-	(1999)
Total under-five deaths (000)	14	(1999)
Neonatal deaths: % of all under-5 deaths	46	(1999)
Neonatal mortality rate (per 1000 live births)	23	(1999)
Infant mortality rate (per 1000 live births)	25	(1999)
SBIR rate (per 1000 total births)	36	(1999)
Total maternal deaths	480	(1999)
Lifetime risk of maternal death (1 in N)	210	(1999)
Total fertility rate (per woman)	4.3	(2010)
Adolescent birth rate (per 1000 women)	92	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for PMCT as a % of total ARTs

Percent HIV+ pregnant women receiving ARTs for PMCT


Uncertainty range around the estimate



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage



DHS 1998-1999

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %) 1 (2008-2009)
Low birthweight incidence (moderate and severe, %) 11 (2008-2009)

Early initiation of breastfeeding (within 1 hr of birth, %) 56 (2008-2009)
Introduction of solid, semi-solid/soft foods (%) -
Vitamin A two dose coverage (%) 36 (2010)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2000 WHO reference population

Exclusive breastfeeding


Percent infants <6 months exclusively breastfed





DEMOGRAPHICS


Total population (000)	9,962	(2009)
Total under-five population (000)	1,606	(2009)
Births (000)	330	(2009)
Birth registration (%)	43	(2009)
Total under-five deaths (000)	48	(2009)
Neonatal deaths: % of all under-5 deaths	81	(2009)
Neonatal mortality rate (per 1000 live births)	30	(2009)
Infant mortality rate (per 1000 live births)	51	(2009)
Skilled birth rate (per 1000 total births)	24	(2009)
Total maternal deaths	2,700	(2009)
Lifetime risk of maternal death (1 in N)	26	(2009)
Total fertility rate (per woman)	5.2	(2009)
Adolescent birth rate (per 1000 women)	158	(2009)



Note: MDG target established by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Skilled attendant at delivery

Percent live births attended by skilled health personnel




Prevention of mother-to-child transmission of HIV

Skilled HIV-prepared women receiving ART for their own health (% of total ART) 27 (2009)



EQUITY


Socioeconomic inequities in coverage



Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization



Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider



NUTRITION

Wasting prevalence (moderate and severe, %)	8	(2006)
Low birthweight incidence (moderate and severe, %)	12	(2008)
Early initiation of breastfeeding (within 1 hr of birth, %)	40	(2006)
Introduction of solid, semi-solid/soft foods (%)	-	-
Vitamin A two dose coverage (%)	87	(2010)

Underweight and stunting prevalence



Exclusive breastfeeding






DEMOGRAPHICS

Total population (000)	1,515	(2009)
Total under-five population (000)	240	(2009)
Births (000)	58	(2009)
Birth registration (%)	34	(2009)
Total under-five deaths (000)	8	(2009)
Neonatal deaths: % of all under-5 deaths	38	(2009)
Neonatal mortality rate (per 1000 live births)	40	(2009)
Infant mortality rate (per 1000 live births)	32	(2009)
SBIR rate (per 1000 total births)	30	(2009)
Total maternal deaths	650	(2009)
Lifetime risk of maternal death (1 in N)	18	(2009)
Total fertility rate (per woman)	5.1	(2009)
Adolescent birth rate (per 1000 women)	187	(2009)

Under-five mortality rate

Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ARTs for PMCT (% of total HIV+)

Percent HIV+ pregnant women receiving ARTs for PMCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



MICS 2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	8	(2010)
Low birthweight incidence (moderate and severe, %)	11	(2010)
Early initiation of breastfeeding (within 1 hr of birth, %)	55	(2010)
Introduction of solid, semi-solid/soft foods (%)	43	(2010)
Vitamin A two dose coverage (%)	100	(2010)

Underweight and stunting prevalence

Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health




DEMOGRAPHICS

Total population (000)	9,988	(2008)
Total under-five population (000)	1,257	(2008)
Births (000)	216	(2008)
Birth registration (%)	81	(2008-2009)
Total under-five deaths (000)	45	(2008)
Neonatal deaths: % of all under-5 deaths	36	(2008)
Neonatal mortality rate (per 1000 live births)	27	(2008)
Infant mortality rate (per 1000 live births)	70	(2008)
Skilled birth rate (per 1000 total births)	15	(2008)
Total maternal deaths	620	(2008)
Lifetime risk of maternal death (1 in N)	98	(2008)
Total fertility rate (per woman)	4.3	(2008)
Adolescent birth rate (per 1000 women)	60	(2008)



MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Skilled attendant at delivery




Prevention of mother-to-child transmission of HIV




EQUITY

Socioeconomic inequities in coverage




CHILD HEALTH

Immunization




Pneumonia treatment



NUTRITION

Wasting prevalence (moderate and severe, %)	30	(2005-2006)
Low birthweight incidence (moderate and severe, %)	25	(2005-2006)
Early initiation of breastfeeding (within 1 hr of birth, %)	44	(2009-2008)
Introduction of solid, semi-solid/soft foods (%)	90	(2009-2008)
Vitamin A two dose coverage (%)	21	(2008)

Underweight and stunting prevalence



Exclusive breastfeeding






DEMOGRAPHICS

Total population (000)	1,224,634	(1999)
Total under-five population (000)	127,979	(2008)
Births (000)	27,310	(2008)
Birth registration (%)	41	(2008, 2009)
Total under-five deaths (000)	1,006	(2008)
Neonatal deaths: % of all under-5 deaths	52	(2008)
Neonatal mortality rate (per 1000 live births)	32	(2008)
Infant mortality rate (per 1000 live births)	48	(2008)
SBIR rate (per 1000 total births)	22	(2008)
Total maternal deaths	63,000	(1999)
Lifetime risk of maternal death (1 in N)	140	(2008)
Total fertility rate (per woman)	2.6	(2008)
Adolescent birth rate (per 1000 women)	43	(2008)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Highly HIV+ pregnant women receiving ART for their own health (% of total HIV+)

Percent HIV+ pregnant women receiving ART for PMCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



NFHS 2005-2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib




Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider

Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)

20 (2005-2006)

Low birthweight incidence (moderate and severe, %)

28 (2005-2006)

Early initiation of breastfeeding (within 1 hr of birth, %)

41 (2007-2008)

Introduction of solid, semi-solid/soft foods (%)


56 (2005-2006)

Vitamin A two dose coverage (%)

34 (2008)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2000 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	239,871	(2009)
Total under-five population (000)	21,579	(2009)
Births (000)	4,372	(2009)
Birth registration (%)	53	(2009)
Total under-five deaths (000)	151	(2009)
Neonatal deaths: % of all under-5 deaths	46	(2009)
Neonatal mortality rate (per 1000 live births)	17	(2009)
Infant mortality rate (per 1000 live births)	27	(2009)
SBIR rate (per 1000 total births)	15	(2009)
Total maternal deaths	30,000	(2009)
Lifetime risk of maternal death (1 in N)	190	(2009)
Total fertility rate (per woman)	2.1	(2009)
Adolescent birth rate (per 1000 women)	52	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV


Eligible HIV+ pregnant women receiving ARTs for PMCT as a % of total HIV+ women (2010)



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage




DHS 2007

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization



Source: WHO/UNICEF

Pneumonia treatment


Percent children <5 years with suspected pneumonia taken to appropriate health provider



NUTRITION

Wasting prevalence (moderate and severe, %)	15	(2007)	Early initiation of breastfeeding (within 1 hr of birth, %)	44	(2007)
Low birthweight incidence (moderate and severe, %)	9	(2007)	Introduction of solid, semi-solid/soft foods (%)	85	(2007)
			Vitamin A two dose coverage (%)	80	(2007)

Underweight and stunting prevalence



Note: Based on 2000 WHO reference population



Exclusive breastfeeding






DEMOGRAPHICS

Total population (000)	31,672	(2009)
Total under five population (000)	5,188	(2009)
Births (000)	1,125	(2009)
Birth registration (%)	95	(2009)
Total under five deaths (000)	44	(2009)
Neonatal deaths (% of all under 5 deaths)	58	(2009)
Neonatal mortality rate (per 1000 live births)	20	(2009)
Infant mortality rate (per 1000 live births)	31	(2009)
Skilled birth rate (per 1000 total births)	9	(2009)
Total maternal deaths	780	(2009)
Lifetime risk of maternal death (1 in N)	300	(2009)
Total fertility rate (per woman)	4.7	(2009)
Adolescent birth rate (per 1000 women)	68	(2009)



MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Skilled attendant at delivery



Prevention of mother-to-child transmission of HIV




EQUITY

Socioeconomic inequities in coverage




CHILD HEALTH

Immunization




Pneumonia treatment



NUTRITION

Wasting prevalence (moderate and severe, %)	6	(2006)
Low birthweight incidence (moderate and severe, %)	15	(2006)
Early initiation of breastfeeding (within 1 hr of birth, %)	31	(2006)
Introduction of solid, semi-solid/soft foods (%)	62	(2006)
Vitamin A two dose coverage (%)	-	-

Underweight and stunting prevalence



Exclusive breastfeeding



Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on care indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

Note: Based on 2006 WHO reference population



DEMOGRAPHICS

Total population (000)	40,511	(1999)
Total under-five population (000)	6,664	(1999)
Births (000)	1,529	(1999)
Birth registration (%)	40	(2008-2009)
Total under-five deaths (000)	122	(1999)
Neonatal deaths: % of all under-5 deaths	35	(1999)
Neonatal mortality rate (per 1000 live births)	28	(1999)
Infant mortality rate (per 1000 live births)	55	(1999)
Stillbirth rate (per 1000 total births)	22	(1999)
Total maternal deaths	7,900	(1999)
Lifetime risk of maternal death (1 in N)	38	(1999)
Total fertility rate (per woman)	4.7	(2009)
Adolescent birth rate (per 1000 women)	106	(1999)

Under-five mortality rate

Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Source: DHS

Prevention of mother-to-child transmission of HIV


Eligible HIV+ pregnant women receiving ARTs for PMCT (% of total HIV+)



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage




DHS 2008-2009

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider




NUTRITION

Wasting prevalence (moderate and severe, %) 7 (2008-2009)
Low birthweight incidence (moderate and severe, %) 8 (2008-2009)

Early initiation of breastfeeding (within 1 hr of birth, %) 58 (2008-2009)
Introduction of solid, semi-solid/soft foods (%) 85 (2008-2009)
Vitamin A two dose coverage (%) 62 (2010)

Underweight and stunting prevalence



Note: Based on 2000 WHO reference population

Exclusive breastfeeding





DEMOGRAPHICS

Total population (000)	24,346	0000
Total under-five population (000)	1,704	0000
Births (000)	348	0000
Birth registration (%)	100	0000
Total under-five deaths (000)	12	0000
Neonatal deaths: % of all under-5 deaths	52	0000
Neonatal mortality rate (per 1000 live births)	18	0000
Infant mortality rate (per 1000 live births)	26	0000
Fertility rate (per 1000 total births)	53	0000
Total maternal deaths	83	0000
Lifetime risk of maternal death (1 in N)	230	0000
Total fertility rate (per woman)	2.0	0000
Adolescent birth rate (per 1000 women)	1	0000

Under-five mortality rate

Deaths per 1000 live births



Source: FAO 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IMBB 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See previous charts for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Skilled HIV+ pregnant women receiving ART for their own health (% of total HIV+)

Percent HIV+ pregnant women receiving ART for PMCT
Uncertainty range around the estimate

No Data

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



No Data


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 5 0000
Low birthweight incidence (moderate and severe, %) 8 0000

Early initiation of breastfeeding (within 1 hr of birth, %) 18 0000
Introduction of solid, semi-solid/soft foods (%) - 0000
Vitamin A two dose coverage (%) 99 0000

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	5,134	(2009)
Total under-five population (000)	595	(2009)
Births (000)	130	(2009)
Birth registration (%)	94	(2009)
Total under-five deaths (000)	5	(2009)
Neonatal deaths: % of all under-5 deaths	46	(2009)
Neonatal mortality rate (per 1000 live births)	29	(2009)
Infant mortality rate (per 1000 live births)	34	(2009)
SBIR* birth rate (per 1000 total births)	36	(2009)
Total maternal deaths	97	(2009)
Lifetime risk of maternal death (1 in N)	150	(2009)
Total fertility rate (per woman)	2.7	(2009)
Adolescent birth rate (per 1000 women)	30	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: UNICEF 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: UNICEF 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV


Eligible HIV+ pregnant women receiving ARTs for PMCT (% of total ARTs)



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage




MICS 2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH


Immunization



Source: WHO/UNICEF

Pneumonia treatment


Percent children <5 years with suspected pneumonia taken to appropriate health provider



NUTRITION

Wasting prevalence (moderate and severe, %)	3	(2006)
Low birthweight incidence (moderate and severe, %)	5	(2006)
Early initiation of breastfeeding (within 1 hr of birth, %)	65	(2006)
Introduction of solid, semi-solid/soft foods (%)	60	(2006)
Vitamin A two dose coverage (%)	97	(2006)

Underweight and stunting prevalence



Note: Based on 2006 WHO reference population


Exclusive breastfeeding





DEMOGRAPHICS


Total population (000)	6,202	(2008)
Total under-five population (000)	983	(2008)
Births (000)	141	(2008)
Birth registration (%)	72	(2008)
Total under-five deaths (000)	8	(2008)
Neonatal deaths (% of all under-5 deaths)	38	(2008)
Neonatal mortality rate (per 1000 live births)	21	(2008)
Infant mortality rate (per 1000 live births)	42	(2008)
Skilled birth rate (per 1000 total births)	54	(2008)
Total maternal deaths	100	(2008)
Lifetime risk of maternal death (1 in N)	69	(2008)
Total fertility rate (per woman)	2.7	(2008)
Adolescent birth rate (per 1000 women)	110	(2008)



Note: MDG target established by Countdown in 2005

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Skilled attendant at delivery

Percent live births attended by skilled health personnel




Prevention of mother-to-child transmission of HIV

Skilled HIV-prepared women receiving ART for their own health (% of total ART)

Percent HIV-prepared women receiving ART for PMCT


Uncertainty range around the estimate



EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	7	(2006)
Low birthweight incidence (moderate and severe, %)	11	(2008)
Early initiation of breastfeeding (within 1 hr of birth, %)	30	(2006)
Introduction of solid, semi-solid/soft foods (%)	41	(2006)
Vitamin A two dose coverage (%)	83	(2010)

Underweight and stunting prevalence

Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





DEMOGRAPHICS

Total population (000)	2,171	(2009)
Total under-five population (000)	274	(2009)
Births (000)	60	(2009)
Birth registration (%)	45	(2009)
Total under-five deaths (000)	5	(2009)
Neonatal deaths: % of all under-5 deaths	42	(2009)
Neonatal mortality rate (per 1000 live births)	35	(2009)
Infant mortality rate (per 1000 live births)	66	(2009)
Stillbirth rate (per 1000 total births)	25	(2009)
Total maternal deaths	310	(2009)
Lifetime risk of maternal death (1 in N)	62	(2009)
Total fertility rate (per woman)	4.2	(2009)
Adolescent birth rate (per 1000 women)	92	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Highly HIV+ pregnant women receiving ARTs for their own health (% of total HIV+) 24 (2010)

Percent HIV+ pregnant women receiving ARTs for PMCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2008

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)	4	(2008)	Early initiation of breastfeeding (within 1 hr of birth, %)	53	(2008)
Low birthweight incidence (moderate and severe, %)	13	(2004)	Introduction of solid, semi-solid/soft foods (%)	68	(2008)
			Vitamin A two dose coverage (%)	38	(2007)

Underweight and stunting prevalence

Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2000 WHO reference population

Exclusive breastfeeding



Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	3,991	(2008)
Total under-five population (000)	961	(2008)
Births (000)	154	(2008)
Birth registration (%)	4	(2008)
Total under-five deaths (000)	15	(2008)
Neonatal deaths: % of all under-5 deaths	34	(2008)
Neonatal mortality rate (per 1000 live births)	34	(2008)
Infant mortality rate (per 1000 live births)	74	(2008)
Stillbirth rate (per 1000 total births)	27	(2008)
Total maternal deaths	1,430	(2008)
Lifetime risk of maternal death (1 in N)	20	(2008)
Total fertility rate (per woman)	5.2	(2008)
Adolescent birth rate (per 1000 women)	177	(2008)



MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS
* See annexes for indicator details


Skilled attendant at delivery

Percent live births attended by skilled health personnel




Prevention of mother-to-child transmission of HIV

Skilled HIV-prepared women receiving ART for their own health (% of total ART) 18 (2010)



EQUITY


Socioeconomic inequities in coverage



DHS 2007
Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.


CHILD HEALTH

Immunization



Source: WHO/UNICEF


Pneumonia treatment



NUTRITION

Wasting prevalence (moderate and severe, %)	3	(2008)
Low birthweight incidence (moderate and severe, %)	14	(2007)
Early initiation of breastfeeding (within 1 hr of birth, %)	44	(2008)
Introduction of solid, semi-solid/soft foods (%)	-	-
Vitamin A two dose coverage (%)	87	(2008)

Underweight and stunting prevalence



Note: Based on 2006 WHO reference population

Exclusive breastfeeding






DEMOGRAPHICS

Total population (000)	20,754	(2008)
Total under-five population (000)	3,306	(2008)
Births (000)	732	(2008)
Birth registration (%)	80	(2008/2009)
Total under-five deaths (000)	44	(2008)
Neonatal deaths: % of all under-5 deaths	37	(2008)
Neonatal mortality rate (per 1000 live births)	22	(2008)
Infant mortality rate (per 1000 live births)	48	(2008)
Stillbirth rate (per 1000 total births)	21	(2008)
Total maternal deaths	3,000	(2008)
Lifetime risk of maternal death (1 in N)	45	(2008)
Total fertility rate (per woman)	4.7	(2008)
Adolescent birth rate (per 1000 women)	147	(2007)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2012

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annex website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Double ARV+ pregnant women receiving AZT for their own health (% of total ARV)


Legend: ■ Percent HIV+ pregnant women receiving AZT for PMCT; ■ Uncertainty range around the estimate



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage



DHS 2008-2009

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

Legend: ■ Percent of children immunized against measles; ■ Percent of children immunized with 3 doses DTP; ■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Legend: ■ Percent children <5 years with suspected pneumonia taken to appropriate health provider; ■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)	95	(2009-2009)
Low birthweight incidence (moderate and severe, %)	38	(2009-2009)
Early initiation of breastfeeding (within 1 hr of birth, %)	72	(2009-2009)
Introduction of solid, semi-solid/soft foods (%)	84	(2009-2009)
Vitamin A two dose coverage (%)	95	(2009)

Underweight and stunting prevalence


Legend: ■ Percent children <5 years who are underweight; ■ Percent children <5 years who are stunted



Note: Based on 2008 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	14,903	(2009)
Total under-five population (000)	2,715	(2009)
Births (000)	663	(2009)
Birth registration (%)	-	(2009)
Total under-five deaths (000)	56	(2009)
Neonatal deaths: % of all under-5 deaths	32	(2009)
Neonatal mortality rate (per 1000 live births)	27	(2009)
Infant mortality rate (per 1000 live births)	58	(2009)
SBIR rate (per 1000 total births)	24	(2009)
Total maternal deaths	3,000	(2009)
Lifetime risk of maternal death (1 in N)	36	(2009)
Total fertility rate (per woman)	6.0	(2010)
Adolescent birth rate (per 1000 women)	1.7	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IAHMs 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Source: DHS 2011

Prevention of mother-to-child transmission of HIV


Eligible HIV+ pregnant women receiving ART for PMCT (% of total HIV+)



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage




DHS 2010

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization



Source: WHO/UNICEF

Pneumonia treatment


Percent children <5 years with suspected pneumonia taken to appropriate health provider



NUTRITION

Wasting prevalence (moderate and severe, %)	4	(2010)	Early initiation of breastfeeding (within 1 hr of birth, %)	58	(2008)
Low birthweight incidence (moderate and severe, %)	13	(2008)	Introduction of solid, semi-solid/soft foods (%)	86	(2010)
			Vitamin A two dose coverage (%)	96	(2010)

Underweight and stunting prevalence



Note: Based on 2000 WHO reference population

Exclusive breastfeeding






DEMOGRAPHICS

Total population (000)	15,370	(2009)
Total under-five population (000)	2,912	(2009)
Births (000)	754	(2009)
Birth registration (%)	81	(2009)
Total under-five deaths (000)	120	(2009)
Neonatal deaths: % of all under-5 deaths	36	(2009)
Neonatal mortality rate (per 1000 live births)	46	(2009)
Infant mortality rate (per 1000 live births)	99	(2009)
SBIR rate (per 1000 total births)	25	(2009)
Total maternal deaths	4,500	(2009)
Lifetime risk of maternal death (1 in 16)	32	(2009)
Total fertility rate (per woman)	6.3	(2010)
Adolescent birth rate (per 1000 women)	100	(2009)

Under-five mortality rate

Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV


Eligible HIV+ pregnant women receiving ART for PMCT (% of total HIV+)



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage




DHS 2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization



Source: WHO/UNICEF


Pneumonia treatment



NUTRITION

Wasting prevalence (moderate and severe, %)	9	(2008)	Early initiation of breastfeeding (within 1 hr of birth, %)	46	(2008)
Low birthweight incidence (moderate and severe, %)	19	(2008)	Introduction of solid, semi-solid/soft foods (%)	25	(2008)
			Vitamin A two dose coverage (%)	99	(2008)

Underweight and stunting prevalence



Note: Based on 2000 WHO reference population

Exclusive breastfeeding






DEMOGRAPHICS

Total population (000)	6,460	(2009)
Total under-five population (000)	513	(2009)
Births (000)	117	(2009)
Birth registration (%)	56	(2009)
Total under-five deaths (000)	13	(2009)
Neonatal deaths: % of all under-5 deaths	36	(2009)
Neonatal mortality rate (per 1000 live births)	39	(2009)
Infant mortality rate (per 1000 live births)	75	(2009)
SBIR rate (per 1000 total births)	27	(2009)
Total maternal deaths	590	(2009)
Lifetime risk of maternal death (1 in N)	41	(2009)
Total fertility rate (per woman)	4.5	(2009)
Adolescent birth rate (per 1000 women)	80	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See annex 2 table for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ARTs for PMCT (% of total HIV+)

54 (2010)

Percent HIV+ pregnant women receiving ARTs for PMCT

Uncertainty range around the estimate




Source: UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



MICS 2007

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNAIDS

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	7	(2010)
Low birthweight incidence (moderate and severe, %)	34	(2007)
Early initiation of breastfeeding (within 1 hr of birth, %)	81	(2010)
Introduction of solid, semi-solid/soft foods (%)	-	-
Vitamin A two dose coverage (%)	87	(2010)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2000 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	118,423	(2009)
Total under-five population (000)	11,005	(2009)
Births (000)	2,217	(2009)
Birth registration (%)	-	
Total under-five deaths (000)	37	(2009)
Neonatal deaths: % of all under-5 deaths	46	(2009)
Neonatal mortality rate (per 1000 live births)	7	(2009)
Infant mortality rate (per 1000 live births)	14	(2009)
SBIR rate (per 1000 total births)	5	(2009)
Total maternal deaths	1,700	(2009)
Lifetime risk of maternal death (1 in N)	503	(2009)
Total fertility rate (per woman)	2.3	(2010)
Adolescent birth rate (per 1000 women)	30	(2009)

Under-five mortality rate

Deaths per 1000 live births



Source: HMN 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Source: HMN 2011

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total HIV+)

Percent HIV+ pregnant women receiving ART for PMCT

Uncertainty range around the estimate



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)	2	(2008)	Early initiation of breastfeeding (within 1 hr of birth, %)	18	(2008)
Low birthweight incidence (moderate and severe, %)	7	(2008)	Introduction of solid, semi-solid/soft foods (%)	-	
			Vitamin A two dose coverage (%)	-	

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2000 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	41,951	(1999)
Total under-five population (000)	8,022	(1999)
Births (000)	623	(1999)
Birth registration (%)	85	(1999)
Total under-five deaths (000)	23	(1999)
Neonatal deaths: % of all under-5 deaths	53	(1999)
Neonatal mortality rate (per 1000 live births)	29	(1999)
Infant mortality rate (per 1000 live births)	30	(1999)
SBIR rate (per 1000 total births)	26	(1999)
Total maternal deaths	720	(1999)
Lifetime risk of maternal death (1 in 16)	360	(1999)
Total fertility rate (per woman)	2.3	(2009)
Adolescent birth rate (per 1000 women)	18	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Source: DHS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total HIV+)

Percent HIV+ pregnant women receiving ART for PMCT


Uncertainty range around the estimate



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage




DHS 2003-2004

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	11	(2003-2004)
Low birthweight incidence (moderate and severe, %)	15	(2003-2004)
Early initiation of breastfeeding (within 1 hr of birth, %)	52	(2003-2004)
Introduction of solid, semi-solid/soft foods (%)	-	-
Vitamin A two dose coverage (%)	63	(2008)

Underweight and stunting prevalence

Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2000 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	23,893	(2008)
Total under-five population (000)	8,876	(2008)
Births (000)	883	(2008)
Birth registration (%)	31	(2008)
Total under-five deaths (000)	114	(2008)
Neonatal deaths: % of all under-5 deaths	30	(2008)
Neonatal mortality rate (per 1000 live births)	30	(2008)
Infant mortality rate (per 1000 live births)	92	(2008)
SBIR rate (per 1000 total births)	26	(2008)
Total maternal deaths	4,800	(2008)
Lifetime risk of maternal death (1 in N)	37	(2008)
Total fertility rate (per woman)	4.9	(2008)
Adolescent birth rate (per 1000 women)	105	(2008)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annexure for indicator definition


Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV


Eligible HIV+ pregnant women receiving ART for PMCT as a % of total HIV+ (2010)



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage




DHS 2008

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization



Source: WHO/UNICEF


Pneumonia treatment



NUTRITION

Wasting prevalence (moderate and severe, %)	4	(2008)	Early initiation of breastfeeding (within 1 hr of birth, %)	83	(2008)
Low birthweight incidence (moderate and severe, %)	18	(2008)	Introduction of solid, semi-solid/soft foods (%)	86	(2008)
			Vitamin A two dose coverage (%)	100	(2010)

Underweight and stunting prevalence



Note: Based on 2000 WHO reference population

Exclusive breastfeeding






DEMOGRAPHICS

Total population (000)	47,964	(1999)
Total under-five population (000)	5,956	(2009)
Births (000)	830	(2009)
Birth registration (%)	72	(2009-2010)
Total under-five deaths (000)	56	(2009)
Neonatal deaths: % of all under-5 deaths	47	(2009)
Neonatal mortality rate (per 1000 live births)	32	(2009)
Infant mortality rate (per 1000 live births)	50	(2009)
Stillbirth rate (per 1000 total births)	26	(2009)
Total maternal deaths	2,400	(2009)
Lifetime risk of maternal death (1 in N)	180	(2009)
Total fertility rate (per woman)	2.0	(2009)
Adolescent birth rate (per 1000 women)	17	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annex 2 table for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Highly HIV+ pregnant women receiving ART for their own health (% of total HIV+)

Percent HIV+ pregnant women receiving ART for PMCT

Uncertainty range around the estimate



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)

Low birthweight incidence (moderate and severe, %)

Early initiation of breastfeeding (within 1 hr of birth, %)

Introduction of solid, semi-solid/soft foods (%)

Vitamin A two dose coverage (%)

Underweight and stunting prevalence

Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2000 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	29,959	(2008)
Total under-five population (000)	8,506	(2008)
Births (000)	724	(2008)
Birth registration (%)	35	(2008)
Total under-five deaths (000)	35	(2008)
Neonatal deaths: % of all under-5 deaths	58	(2008)
Neonatal mortality rate (per 1000 live births)	28	(2008)
Infant mortality rate (per 1000 live births)	41	(2008)
SBIR rate (per 1000 total births)	23	(2008)
Total maternal deaths	2,800	(2008)
Lifetime risk of maternal death (1 in N)	80	(2008)
Total fertility rate (per woman)	2.7	(2008)
Adolescent birth rate (per 1000 women)	106	(2008)

Under-five mortality rate

Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total HIV+)

Percent HIV+ pregnant women receiving ART for PMCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2006
Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics



Source: DHS, MICS, Other NS

NUTRITION

Wasting prevalence (moderate and severe, %)	13	(2008)	Early initiation of breastfeeding (within 1 hr of birth, %)	35	(2008)
Low birthweight incidence (moderate and severe, %)	21	(2008)	Introduction of solid, semi-solid/soft foods (%)	71	(2008)
			Vitamin A two dose coverage (%)	81	(2008)

Underweight and stunting prevalence

Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2000 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	15,512	(2009)
Total under-five population (000)	3,085	(2009)
Births (000)	750	(2009)
Birth registration (%)	32	(2009)
Total under-five deaths (000)	100	(2009)
Neonatal deaths: % of all under-5 deaths	34	(2009)
Neonatal mortality rate (per 1000 live births)	32	(2009)
Infant mortality rate (per 1000 live births)	73	(2009)
SBIR rate (per 1000 total births)	25	(2009)
Total maternal deaths	6,500	(2009)
Lifetime risk of maternal death (1 in N)	16	(2009)
Total fertility rate (per woman)	7.1	(2009)
Adolescent birth rate (per 1000 women)	209	(2009)

Under-five mortality rate

Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total HIV+)

Percent HIV+ pregnant women receiving ART for PMCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS/MICS 2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	15	(2010)	Early initiation of breastfeeding (within 1 hr of birth, %)	42	(2010)
Low birthweight incidence (moderate and severe, %)	27	(2008)	Introduction of solid, semi-solid/soft foods (%)	-	-
			Vitamin A two dose coverage (%)	88	(2010)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	156,423	(2009)
Total under-five population (000)	26,340	(2009)
Births (000)	6,332	(2009)
Birth registration (%)	30	(2009)
Total under-five deaths (000)	861	(2009)
Neonatal deaths: % of all under-5 deaths	29	(2009)
Neonatal mortality rate (per 1000 live births)	40	(2009)
Infant mortality rate (per 1000 live births)	88	(2009)
SBIR rate (per 1000 total births)	42	(2009)
Total maternal deaths	50,000	(2009)
Lifetime risk of maternal death (1 in N)	33	(2009)
Total fertility rate (per woman)	5.5	(2009)
Adolescent birth rate (per 1000 women)	123	(2009)

Under-five mortality rate

Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total HIV+)

Percent HIV+ pregnant women receiving ART for PMCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2008

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	14	(2008)	Early initiation of breastfeeding (within 1 hr of birth, %)	38	(2008)
Low birthweight incidence (moderate and severe, %)	12	(2008)	Introduction of solid, semi-solid/soft foods (%)	76	(2008)
			Vitamin A two dose coverage (%)	81	(2008)

Underweight and stunting prevalence

Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2008 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	173,584	(2009)
Total under-five population (000)	21,426	(2009)
Births (000)	4,741	(2009)
Birth registration (%)	27	(2006-2007)
Total under-five deaths (000)	423	(2009)
Neonatal deaths: % of all under-5 deaths	46	(2009)
Neonatal mortality rate (per 1000 live births)	41	(2009)
Infant mortality rate (per 1000 live births)	76	(2009)
SBIR rate (per 1000 total births)	47	(2009)
Total maternal deaths	14,000	(2009)
Lifetime risk of maternal death (1 in %)	90	(2009)
Total fertility rate (per woman)	4.4	(2009)
Adolescent birth rate (per 1000 women)	16	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV


Eligible HIV+ pregnant women receiving ARTs for PMCT (% of total HIV+)



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage




DHS 2006-2007

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization



Source: WHO/UNICEF


Pneumonia treatment



NUTRITION

Wasting prevalence (moderate and severe, %)	14	(2007)
Low birthweight incidence (moderate and severe, %)	32	(2006-2007)
Early initiation of breastfeeding (within 1 hr of birth, %)	29	(2006-2007)
Introduction of solid, semi-solid/soft foods (%)	-	
Vitamin A two dose coverage (%)	87	(2008)

Underweight and stunting prevalence



Note: Based on 2000 WHO reference population

Exclusive breastfeeding






DEMOGRAPHICS

Total population (000)	6,358	(2008)
Total under-five population (000)	962	(2008)
Births (000)	207	(2008)
Birth registration (%)	-	-
Total under-five deaths (000)	12	(2008)
Neonatal deaths: % of all under-5 deaths	39	(2008)
Neonatal mortality rate (per 1000 live births)	23	(2008)
Infant mortality rate (per 1000 live births)	47	(2008)
Stillbirth rate (per 1000 total births)	15	(2008)
Total maternal deaths	530	(2008)
Lifetime risk of maternal death (1 in N)	96	(2008)
Total fertility rate (per woman)	4.8	(2008)
Adolescent birth rate (per 1000 women)	70	(2008)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annex table for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Source: DHS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for PMCT (% of total HIV+)

Percent HIV+ pregnant women receiving ART for PMCT

Uncertainty range around the estimate



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics




Source: DHS

NUTRITION

Wasting prevalence (moderate and severe, %)	4	(2008)	Early initiation of breastfeeding (within 1 hr of birth, %)	-	(2008)
Low birthweight incidence (moderate and severe, %)	11	(2008)	Introduction of solid, semi-solid/soft foods (%)	-	(2008)
			Vitamin A two dose coverage (%)	14	(2008)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2000 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	29,077	(2008)
Total under-five population (000)	2,900	(2008)
Births (000)	594	(2008)
Birth registration (%)	92	(2008)
Total under-five deaths (000)	11	(2008)
Neonatal deaths: % of all under-5 deaths	49	(2008)
Neonatal mortality rate (per 1000 live births)	9	(2008)
Infant mortality rate (per 1000 live births)	15	(2008)
SBIR* birth rate (per 1000 total births)	30	(2008)
Total maternal deaths	600	(2008)
Lifetime risk of maternal death (1 in N)	170	(2008)
Total fertility rate (per woman)	2.5	(2008)
Adolescent birth rate (per 1000 women)	10	(2008)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ARTs for PMCT (% of total HIV+)

Percent HIV+ pregnant women receiving ARTs for PMCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2004

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)

Low birthweight incidence (moderate and severe, %)

Early initiation of breastfeeding (within 1 hr of birth, %)

Introduction of solid, semi-solid/soft foods (%)

Vitamin A two dose coverage (%)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2000 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	94,263	(2008)
Total under-five population (000)	11,254	(2008)
Births (000)	2,344	(2008)
Birth registration (%)	83	(2008)
Total under-five deaths (000)	66	(2008)
Neonatal deaths: % of all under-5 deaths	48	(2008)
Neonatal mortality rate (per 1000 live births)	24	(2008)
Infant mortality rate (per 1000 live births)	23	(2008)
SBIR rate (per 1000 total births)	36	(2008)
Total maternal deaths	2,100	(2008)
Lifetime risk of maternal death (1 in N)	330	(2008)
Total fertility rate (per woman)	3.1	(2008)
Adolescent birth rate (per 1000 women)	33	(2008)

Under-five mortality rate

Deaths per 1000 live births



Source: UNICEF 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: ADB/WHO 2009

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Source: UNICEF/UNAIDS/WHO

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total HIV+)

Percent HIV+ pregnant women receiving ART for PMCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2008

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics



Source: DHS

NUTRITION

Wasting prevalence (moderate and severe, %)	7	(2008)	Early initiation of breastfeeding (within 1 hr of birth, %)	54	(2008)
Low birthweight incidence (moderate and severe, %)	21	(2008)	Introduction of solid, semi-solid/soft foods (%)	90	(2008)
			Vitamin A two dose coverage (%)	81	(2008)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2008 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Source: DHS



DEMOGRAPHICS

Total population (000)	10,624	(2009)
Total under-five population (000)	1,831	(2009)
Births (000)	430	(2009)
Birth registration (%)	82	(2009)
Total under-five deaths (000)	38	(2009)
Neonatal deaths: % of all under-5 deaths	34	(2009)
Neonatal mortality rate (per 1000 live births)	29	(2009)
Infant mortality rate (per 1000 live births)	50	(2009)
SBIR rate (per 1000 total births)	23	(2009)
Total maternal deaths	2,200	(2009)
Lifetime risk of maternal death (1 in N)	35	(2009)
Total fertility rate (per woman)	3.4	(2009)
Adolescent birth rate (per 1000 women)	43	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for PMCT (% of total ARTs)

Percent HIV+ pregnant women receiving ARTs for PMCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2005

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)

Low birthweight incidence (moderate and severe, %)


Early initiation of breastfeeding (within 1 hr of birth, %)

Introduction of solid, semi-solid/soft foods (%)

Vitamin A two dose coverage (%)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2000 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	245	(2008)
Total under-five population (000)	23	(2008)
Births (000)	5	(2008)
Birth registration (%)	75	(2008-2009)
Total under-five deaths (000)	0	(2008)
Neonatal deaths: % of all under-5 deaths	33	(2008)
Neonatal mortality rate (per 1000 live births)	25	(2008)
Infant mortality rate (per 1000 live births)	54	(2008)
Stillbirth rate (per 1000 total births)	22	(2008)
Total maternal deaths	-	-
Lifetime risk of maternal death (1 in N)	-	-
Total fertility rate (per woman)	4.7	(2008)
Adolescent birth rate (per 1000 women)	130	(2008)

Under-five mortality rate

Deaths per 1000 live births



Source: IHME 2011


Maternal mortality ratio

Deaths per 100,000 live births

No Data

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See annex table for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Prevention of mother-to-child transmission of HIV

Highly HIV+ pregnant women receiving ARTs for their own health (% of total HIV+)

Percent HIV+ pregnant women receiving ARTs for PMCT


Uncertainty range around the estimate



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage



DHS 2008-2009

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	11	(2008-2009)
Low birthweight incidence (moderate and severe, %)	8	(2008)
Early initiation of breastfeeding (within 1 hr of birth, %)	45	(2008-2009)
Introduction of solid, semi-solid/soft foods (%)	74	(2008-2009)
Vitamin A two dose coverage (%)	81	(2008)

Underweight and stunting prevalence

Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	12,434	(2008)
Total under-five population (000)	2,081	(2008)
Births (000)	405	(2008)
Birth registration (%)	50	(2008)
Total under-five deaths (000)	34	(2008)
Neonatal deaths: % of all under-5 deaths	37	(2008)
Neonatal mortality rate (per 1000 live births)	27	(2008)
Infant mortality rate (per 1000 live births)	50	(2008)
Stillbirth rate (per 1000 total births)	34	(2008)
Total maternal deaths	1,000	(2008)
Lifetime risk of maternal death (1 in N)	46	(2008)
Total fertility rate (per woman)	4.8	(2008)
Adolescent birth rate (per 1000 women)	96	(2008)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for PMCT (% of total HIV+)

Percent HIV+ pregnant women receiving ART for PMCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2005

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib




Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider

Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)	9	(2008)	Early initiation of breastfeeding (within 1 hr of birth, %)	23	(2008)
Low birthweight incidence (moderate and severe, %)	15	(2008)	Introduction of solid, semi-solid/soft foods (%)	-	-
			Vitamin A two dose coverage (%)	87	(2008)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2000 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	5,968	(2008)
Total under-five population (000)	970	(2008)
Births (000)	226	(2008)
Birth registration (%)	51	(2008)
Total under-five deaths (000)	28	(2008)
Neonatal deaths: % of all under-5 deaths	26	(2008)
Neonatal mortality rate (per 1000 live births)	45	(2008)
Infant mortality rate (per 1000 live births)	114	(2008)
Fertility rate (per 1000 total births)	30	(2008)
Total maternal deaths	2,200	(2008)
Relative risk of maternal death (1 in N)	71	(2008)
Total fertility rate (per woman)	5.0	(2008)
Adolescent birth rate (per 1000 women)	143	(2008)

Under-five mortality rate

Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Highly HIV+ pregnant women receiving ARTs for own health (% of total HIV+)

Percent HIV+ pregnant women receiving ARTs for PMCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2008

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)	11	(2008)	Early initiation of breastfeeding (within 1 hr of birth, %)	51	(2008)
Low birthweight incidence (moderate and severe, %)	14	(2008)	Introduction of solid, semi-solid/soft foods (%)	78	(2008)
			Vitamin A two dose coverage (%)	100	(2008)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2000 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	546	(2009)
Total under-five population (000)	90	(2009)
Births (000)	17	(2009)
Birth registration (%)	80	(2009)
Total under-five deaths (000)	0	(2009)
Neonatal deaths: % of all under-5 deaths	47	(2009)
Neonatal mortality rate (per 1000 live births)	12	(2009)
Infant mortality rate (per 1000 live births)	23	(2009)
Stillbirth rate (per 1000 total births)	15	(2009)
Total maternal deaths	16	(2009)
Lifetime risk of maternal death (1 in N)	280	(2009)
Total fertility rate (per woman)	4.2	(2009)
Adolescent birth rate (per 1000 women)	70	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Highly HIV+ pregnant women receiving ART for their own health (% of total HIV+)

Percent HIV+ pregnant women receiving ART for PMCT

Uncertainty range around the estimate

No Data

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib




Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider

Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)

4 (2007)

Low birthweight incidence (moderate and severe, %)

13 (2007)

Early initiation of breastfeeding (within 1 hr of birth, %)


75 (2007)

Introduction of solid, semi-solid/soft foods (%)

Vitamin A two dose coverage (%)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2000 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





DEMOGRAPHICS

Total population (000)	9,333	(2009)
Total under-five population (000)	1,667	(2009)
Births (000)	408	(2009)
Birth registration (%)	2	(2009)
Total under-five deaths (000)	70	(2009)
Neonatal deaths: % of all under-5 deaths	30	(2009)
Neonatal mortality rate (per 1000 live births)	32	(2009)
Infant mortality rate (per 1000 live births)	106	(2009)
Stillbirth rate (per 1000 total births)	30	(2009)
Total maternal deaths	4,800	(2009)
Lifetime risk of maternal death (1 in N)	14	(2009)
Total fertility rate (per woman)	6.3	(2009)
Adolescent birth rate (per 1000 women)	123	(2009)

Under-five mortality rate

Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annexes for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ARTs for non-breast (% of total ARTs) (2010)

Percent HIV+ pregnant women receiving ARTs for PMCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



MICS 2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	13	(2006)	Early initiation of breastfeeding (within 1 hr of birth, %)	26	(2006)
Low birthweight incidence (moderate and severe, %)	1	(2006)	Introduction of solid, semi-solid/soft foods (%)	18	(2006)
			Vitamin A two dose coverage (%)	62	(2006)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





DEMOGRAPHICS

Total population (000)	50,884	(2008)
Total under-five population (000)	5,041	(2008)
Births (000)	1,059	(2008)
Birth registration (%)	92	(2008)
Total under-five deaths (000)	58	(2008)
Neonatal deaths: % of all under-5 deaths	32	(2008)
Neonatal mortality rate (per 1000 live births)	28	(2008)
Infant mortality rate (per 1000 live births)	41	(2008)
Stillbirth rate (per 1000 total births)	20	(2008)
Total maternal deaths	4,500	(2008)
Lifetime risk of maternal death (1 in N)	108	(2008)
Total fertility rate (per woman)	2.5	(2008)
Adolescent birth rate (per 1000 women)	34	(2008)

Under-five mortality rate

Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Source: DHS

Prevention of mother-to-child transmission of HIV


Highly HIV+ pregnant women receiving ARTs for PMCT: 12 (2010)



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage




DHS 2003

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization



Source: WHO/UNICEF

Pneumonia treatment



NUTRITION

Wasting prevalence (moderate and severe, %)	5	(2008)	Early initiation of breastfeeding (within 1 hr of birth, %)	81	(2008)
Low birthweight incidence (moderate and severe, %)	15	(2008)	Introduction of solid, semi-solid/soft foods (%)	-	-
			Vitamin A two dose coverage (%)	89	(2008)

Underweight and stunting prevalence



Note: Based on 2008 WHO reference population

Exclusive breastfeeding






DEMOGRAPHICS

Total population (000)	1,186	(2009)
Total under-five population (000)	157	(2009)
Births (000)	25	(2009)
Birth registration (%)	80	(2009, 2010)
Total under-five deaths (000)	3	(2009)
Neonatal deaths: % of all under-5 deaths	29	(2009)
Neonatal mortality rate (per 1000 live births)	21	(2009)
Infant mortality rate (per 1000 live births)	50	(2009)
Stillbirth rate (per 1000 total births)	18	(2009)
Total maternal deaths	150	(2009)
Lifetime risk of maternal death (1 in N)	75	(2009)
Total fertility rate (per woman)	3.4	(2010)
Adolescent birth rate (per 1000 women)	111	(2009)

Under-five mortality rate

Deaths per 1000 live births



Source: IHME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2012

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* Not applicable for selected datasets

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Highly HIV+ pregnant women receiving ARTs for their own health (% of total ARTs) (1: 2010)




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2006-2007

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

— Percent of children immunized against measles
— Percent of children immunized with 3 doses DTP
— Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 1 (2008)
Low birthweight incidence (moderate and severe, %) 9 (2008)

Early initiation of breastfeeding (within 1 hr of birth, %) 64 (2008)
Introduction of solid, semi-solid/soft foods (%) - -
Vitamin A two dose coverage (%) 38 (2008)

Underweight and stunting prevalence

■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2000 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	6,879	(2009)
Total under-five population (000)	871	(2009)
Births (000)	192	(2009)
Birth registration (%)	88	(2009)
Total under-five deaths (000)	12	(2009)
Neonatal deaths: % of all under-5 deaths	41	(2009)
Neonatal mortality rate (per 1000 live births)	25	(2009)
Infant mortality rate (per 1000 live births)	52	(2009)
Stillbirth rate (per 1000 total births)	12	(2009)
Total maternal deaths	120	(2009)
Lifetime risk of maternal death (1 in N)	483	(2009)
Total fertility rate (per woman)	3.3	(2009)
Adolescent birth rate (per 1000 women)	27	(2009)

Under-five mortality rate

Deaths per 1000 live births



Source: UNICEF 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: UNICEF 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ARTs for PMCT 41 (2010)

Percent HIV+ pregnant women receiving ARTs for PMCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



MICS 2005

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	7	(2008)	Early initiation of breastfeeding (within 1 hr of birth, %)	57	(2008)
Low birthweight incidence (moderate and severe, %)	10	(2008)	Introduction of solid, semi-solid/soft foods (%)	-	-
			Vitamin A two dose coverage (%)	95	(2008)

Underweight and stunting prevalence

Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2000 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	44,843	(1999)
Total under-five population (000)	8,030	(1999)
Births (000)	1,862	(1999)
Birth registration (%)	16	(1999)
Total under-five deaths (000)	133	(1999)
Neonatal deaths: % of all under-5 deaths	36	(1999)
Neonatal mortality rate (per 1000 live births)	26	(1999)
Infant mortality rate (per 1000 live births)	50	(1999)
SBIR rate (per 1000 total births)	26	(1999)
Total maternal deaths	14,000	(1999)
Lifetime risk of maternal death (1 in N)	33	(1999)
Total fertility rate (per woman)	5.5	(2010)
Adolescent birth rate (per 1000 women)	118	(2009)

Under-five mortality rate

Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Source: DHS

Prevention of mother-to-child transmission of HIV

Highly HIV+ pregnant women receiving ARTs for their own health (% of total HIV+)




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2010

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 5 (2010)
Low birthweight incidence (moderate and severe, %) 22 (2004-2005)

Early initiation of breastfeeding (within 1 hr of birth, %) 49 (2010)
Introduction of solid, semi-solid/soft foods (%) 82 (2010)
Vitamin A two dose coverage (%) 99 (2010)

Underweight and stunting prevalence

■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2000 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	6,028	(2009)
Total under-five population (000)	864	(2009)
Births (000)	133	(2009)
Birth registration (%)	72	(2009)
Total under-five deaths (000)	19	(2009)
Neonatal deaths: % of all under-5 deaths	32	(2009)
Neonatal mortality rate (per 1000 live births)	32	(2009)
Infant mortality rate (per 1000 live births)	66	(2009)
Stillbirth rate (per 1000 total births)	25	(2009)
Total maternal deaths	740	(2009)
Lifetime risk of maternal death (1 in 16)	67	(2009)
Total fertility rate (per woman)	4.1	(2010)
Adolescent birth rate (per 1000 women)	89	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Source: DHS, MICS, Other NS, pMICS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ARTs for their own health (% of total ARTs) 24 (2010)

Percent HIV+ pregnant women receiving ARTs for PMCT




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



MICS 2006


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization

— Percent of children immunized against measles
— Percent of children immunized with 3 doses DTP
— Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)	5	(2011)	Early initiation of breastfeeding (within 1 hr of birth, %)	36	(2008)
Low birthweight incidence (moderate and severe, %)	11	(2010)	Introduction of solid, semi-solid/soft foods (%)	44	(2010)
			Vitamin A two dose coverage (%)	100	(2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2000 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	5,042	(2009)
Total under-five population (000)	506	(2009)
Births (000)	109	(2009)
Birth registration (%)	96	(2009)
Total under-five deaths (000)	6	(2009)
Neonatal deaths: % of all under-5 deaths	44	(2009)
Neonatal mortality rate (per 1000 live births)	23	(2009)
Infant mortality rate (per 1000 live births)	47	(2009)
Stillbirth rate (per 1000 total births)	13	(2009)
Total maternal deaths	35	(2009)
Lifetime risk of maternal death (1 in N)	500	(2009)
Total fertility rate (per woman)	2.4	(2009)
Adolescent birth rate (per 1000 women)	21	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: UNICEF 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IAHG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total HIV+)

Percent HIV+ pregnant women receiving ART for PMCT


Uncertainty range around the estimate

No Data

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider

Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %) 7 (2006)
Low birthweight incidence (moderate and severe, %) 4 (2006)


Early initiation of breastfeeding (within 1 hr of birth, %) 60 (2006)

Introduction of solid, semi-solid/soft foods (%) -

Vitamin A two dose coverage (%) -

Underweight and stunting prevalence

Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2000 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	33,425	(2009)
Total under-five population (000)	6,465	(2009)
Births (000)	1,554	(2009)
Birth registration (%)	21	(2009)
Total under-five deaths (000)	141	(2009)
Neonatal deaths: % of all under-5 deaths	38	(2009)
Neonatal mortality rate (per 1000 live births)	26	(2009)
Infant mortality rate (per 1000 live births)	63	(2009)
Stillbirth rate (per 1000 total births)	20	(2009)
Total maternal deaths	6,300	(2009)
Lifetime risk of maternal death (1 in 16)	35	(2009)
Total fertility rate (per woman)	6.1	(2009)
Adolescent birth rate (per 1000 women)	139	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total HIV+)

Percent HIV+ pregnant women receiving ART for PMCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	8	(2008)	Early initiation of breastfeeding (within 1 hr of birth, %)	42	(2008)
Low birthweight incidence (moderate and severe, %)	14	(2008)	Introduction of solid, semi-solid/soft foods (%)	75	(2008)
			Vitamin A two dose coverage (%)	64	(2008)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2008 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	27,445	(2009)
Total under-five population (000)	2,736	(2009)
Births (000)	567	(2009)
Birth registration (%)	100	(2009)
Total under-five deaths (000)	11	(2009)
Neonatal deaths: % of all under-5 deaths	43	(2009)
Neonatal mortality rate (per 1000 live births)	23	(2009)
Infant mortality rate (per 1000 live births)	44	(2009)
Stillbirth rate (per 1000 total births)	6	(2009)
Total maternal deaths	170	(2009)
Lifetime risk of maternal death (1 in N)	1,600	(2009)
Total fertility rate (per woman)	2.4	(2009)
Adolescent birth rate (per 1000 women)	26	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: UNICEF 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IABR 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for PMCT (% of total HIV+)

Percent HIV+ pregnant women receiving ART for PMCT

Unertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




MICS 2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	5	(2008)
Low birthweight incidence (moderate and severe, %)	5	(2008)
Early initiation of breastfeeding (within 1 hr of birth, %)	67	(2008)
Introduction of solid, semi-solid/soft foods (%)	47	(2008)
Vitamin A two dose coverage (%)	94	(2008)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2008 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	87,848	(2008)
Total under-five population (000)	7,186	(2008)
Births (000)	1,467	(2008)
Birth registration (%)	88	(2008)
Total under-five deaths (000)	34	(2008)
Neonatal deaths: % of all under-5 deaths	52	(2008)
Neonatal mortality rate (per 1000 live births)	12	(2008)
Infant mortality rate (per 1000 live births)	29	(2008)
SBIR* birth rate (per 1000 total births)	15	(2008)
Total maternal deaths	640	(2008)
Lifetime risk of maternal death (1 in N)	950	(2008)
Total fertility rate (per woman)	1.8	(2008)
Adolescent birth rate (per 1000 women)	35	(2008)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See annex 2 table for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV


Eligible HIV+ pregnant women receiving ART for PMCT (% of total HIV+)



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage




MICS 2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization



Source: WHO/UNICEF

Pneumonia treatment


Percent children <5 years with suspected pneumonia taken to appropriate health provider



NUTRITION

Wasting prevalence (moderate and severe, %)	10	(2008)
Low birthweight incidence (moderate and severe, %)	5	(2008)
Early initiation of breastfeeding (within 1 hr of birth, %)	58	(2008)
Introduction of solid, semi-solid/soft foods (%)	78	(2008)
Vitamin A two dose coverage (%)	95	(2008)

Underweight and stunting prevalence



Note: Based on 2008 WHO reference population

Exclusive breastfeeding






DEMOGRAPHICS

Total population (000)	24,054	(2009)
Total under-five population (000)	4,057	(2009)
Births (000)	929	(2009)
Birth registration (%)	22	(2009)
Total under-five deaths (000)	68	(2009)
Neonatal deaths: % of all under-5 deaths	43	(2009)
Neonatal mortality rate (per 1000 live births)	32	(2009)
Infant mortality rate (per 1000 live births)	57	(2009)
SBIR rate (per 1000 total births)	25	(2009)
Total maternal deaths	1,800	(2009)
Lifetime risk of maternal death (1 in N)	91	(2009)
Total fertility rate (per woman)	5.2	(2009)
Adolescent birth rate (per 1000 women)	80	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annexes for indicator definitions

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Source: UNICEF/WHO/UNFPA

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ARTs for their own health (% of total HIV+)

Percent HIV+ pregnant women receiving ARTs for PMCT

Uncertainty range around the estimate




Source: UNICEF/WHO/UNFPA

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



MICS 2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	15	(2005)
Low birthweight incidence (moderate and severe, %)	12	(2005)
Early initiation of breastfeeding (within 1 hr of birth, %)	30	(2006)
Introduction of solid, semi-solid/soft foods (%)	-	-
Vitamin A two dose coverage (%)	47	(2006)

Underweight and stunting prevalence

Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	14,089	(2009)
Total under-five population (000)	2,412	(2009)
Births (000)	600	(2009)
Birth registration (%)	14	(2009)
Total under-five deaths (000)	60	(2009)
Neonatal deaths: % of all under-5 deaths	30	(2009)
Neonatal mortality rate (per 1000 live births)	30	(2009)
Infant mortality rate (per 1000 live births)	68	(2009)
SBIR* birth rate (per 1000 total births)	26	(2009)
Total maternal deaths	2,600	(2009)
Lifetime risk of maternal death (1 in %)	38	(2009)
Total fertility rate (per woman)	6.3	(2009)
Adolescent birth rate (per 1000 women)	151	(2009)

Under-five mortality rate

Deaths per 1000 live births



Source: IHME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: IHME 2011

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See annexure for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV


Highly HIV+ pregnant women receiving ARTs for PMCT (% of total HIV+)



Source: UNICEF/UNAIDS/WHO

EQUITY


Socioeconomic inequities in coverage



DHS 2007
Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH


Immunization



Source: WHO/UNICEF

Pneumonia treatment


Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)	8	(2007)	Early initiation of breastfeeding (within 1 hr of birth, %)	57	(2007)
Low birthweight incidence (moderate and severe, %)	11	(2007)	Introduction of solid, semi-solid/soft foods (%)	94	(2007)
			Vitamin A two dose coverage (%)	82	(2007)

Underweight and stunting prevalence



Note: Based on 2000 WHO reference population


Exclusive breastfeeding






DEMOGRAPHICS

Total population (000)	12,571	(2009)
Total under-five population (000)	1,602	(2009)
Births (000)	374	(2009)
Birth registration (%)	38	(2009)
Total under-five deaths (000)	29	(2009)
Neonatal deaths: % of all under-5 deaths	35	(2009)
Neonatal mortality rate (per 1000 live births)	27	(2009)
Infant mortality rate (per 1000 live births)	51	(2009)
Stillbirth rate (per 1000 total births)	20	(2009)
Total maternal deaths	3,000	(2009)
Lifetime risk of maternal death (1 in N)	42	(2009)
Total fertility rate (per woman)	3.3	(2009)
Adolescent birth rate (per 1000 women)	101	(2009)



MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Skilled attendant at delivery




Prevention of mother-to-child transmission of HIV




EQUITY

Socioeconomic inequities in coverage




CHILD HEALTH

Immunization




Pneumonia treatment




NUTRITION

Wasting prevalence (moderate and severe, %)	3 (2007-2011)	Early initiation of breastfeeding (within 1 hr of birth, %)	69 (2008-2008)
Low birthweight incidence (moderate and severe, %)	11 (2008-2008)	Introduction of solid, semi-solid/soft foods (%)	76 (2008-2008)
		Vitamin A two dose coverage (%)	89 (2010)

Underweight and stunting prevalence



Exclusive breastfeeding



Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequity. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

Notes: Based on 2008 WHO reference population

Each year, **millions** of women and children die in pregnancy, childbirth and from preventable disease.

These are not statistics. They are people with names and faces.



**EVERY WOMAN
EVERY CHILD**



These deaths can and must be stopped. We know what works.

Through the *Every Woman Every Child* effort, the **United Nations** and its partners are working to save the lives of 16 million women and children by 2015.

www.everywomaneverychild.org

Together we can empower women around the world to be healthy and safe and to protect their children from preventable illness.